

General elements in anticanonical systems of threefolds with divisorial contractions and applications to classification

Masayuki Kawakita

Abstract

We treat threefolds with divisorial contractions whose exceptional divisors contract to compound Du Val points. We prove that general elements in their anticanonical systems around the exceptional divisors have at worst Du Val singularities. As applications to classification, we describe divisorial contractions to compound A_n points, and moreover we conclude that discrepancies of divisorial contractions to compound D_n or E_n points are at most four.

1 Introduction

This paper aims at completion of the explicit study of three dimensional divisorial contractions whose exceptional divisors contract to compound Du Val points after [Ka1] and [Ka2].

It is M. Reid who has pointed out that general elements in the anticanonical systems of threefolds have at worst Du Val singularities under appropriate situations arising from contractions of extremal faces, and it has pervaded us as the general elephant conjecture. [Sh], [Re2] and [Ta] sustain it by affirmative answers for Fano threefolds with singularities and this approach settles the existence problem of three dimensional flips [Mo2][KM1].

Returning to our case of divisorial contractions, let $f: (Y \supset E) \rightarrow (X \ni P)$ be a germ of a three dimensional divisorial contraction whose exceptional divisor E contracts to a point P . His conjecture claims that general elements in the anticanonical system $|-K_Y|$ of Y around E have Du Val singularities only. This statement is analogue to the flipping case and we can start at a similar stage once we take the intersection C of E and the strict transform of a general hyperplane section on X , because the first cohomology of C vanishes.

There lies, however, a crucial difference in C between our case and the flipping case. For any flipping curve in a threefold the first cohomology of the structure sheaf of any closed subscheme supported on this curve always vanishes, but it does not hold for our curve C . Nevertheless provided that P

is Gorenstein, or equivalently, compound Du Val, numerical information on f in [Ka1] benefits analysis of the local structure of $C \subset Y$ through delicate attention to the way of local embedding of Y into the tangent space at each non-Gorenstein point. It stores the behaviour of global sections in $|-K_Y|$, and finally concludes the existence of Du Val sections (2.7).

Theorem 1.1. *Let $f: (Y \supset E) \rightarrow (X \ni P)$ be a germ of a divisorial contraction whose exceptional divisor E contracts to a cDV point P . Then a general element in $|-K_Y|$ has at worst Du Val singularities.*

Let S be a surface on Y defined by a general element in $|-K_Y|$ and let S_X be its strict transform on X . Our main theorem guarantees that S and S_X have Du Val singularities only and that the induced morphism $S \rightarrow S_X$ factors the minimal resolution of S_X . We show it together with providing information on the partial resolution $S \rightarrow S_X$ (2.8). In some special cases, to be precise in types O and I in (3.2), we see immediately that a general hyperplane section S_X on X gives a Du Val section S . Considerably delicate analysis is required in the remaining cases, but thanks to this analysis we can obtain the strong version of the general elephant conjecture, which asserts that the type of a Du Val singularity $Q \in S$ for any point Q is nothing but that of a general Du Val section of a germ $Q \in Y$. In most cases it is proved by the direct search for S , whereas in some exceptional cases where $-K_Y$ is linearly equivalent to the sum of E and a Cartier divisor L we obtain a desired S by showing that E has a Du Val singularity of the required type at any $Q \in S$ and that L has no base points. We add that complete description of the partial resolution $S \rightarrow S_X$ is also obtained. For instance, its exceptional locus is irreducible in nearly every case.

We apply this theorem to classification of divisorial contractions naturally. We describe these contractions to cA_n points (2.12), following smooth and cA_1 case [Ka1][Ka2].

Theorem 1.2. *Assume that P is cA_n ($n \geq 2$). Then one of the following holds.*

- (i) *Under a suitable identification $P \in X \cong o \in (x_1x_2 + g(x_3, x_4) = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4$ -space, f is the weighted blowup with weights $\text{wt}(x_1, x_2, x_3, x_4) = (r_1, r_2, a, 1)$, where a divides $r_1 + r_2$ and is coprime to r_1 and r_2 , the weighted order of g with weights $\text{wt}(x_3, x_4) = (a, 1)$ is $r_1 + r_2$, and the coefficient of $x_3^{(r_1+r_2)/a}$ in g is not zero. Moreover any such f is a divisorial contraction.*
- (ii) *P is a cA_2 point isomorphic to $o \in (x_1x_2 + x_3^3 + g_{\geq 4}(x_3, x_4) = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4$ -space, where the total order of $g_{\geq 4}$ with respect to x_3, x_4 is greater than or equal to 4, Y has exactly one non-Gorenstein point Q , which is isomorphic to $o \in (y_1^2 + y_2^2 + y_3^2 + y_4^3 = 0)$ in the quotient space of $\mathbb{C}^4 = y_1y_2y_3y_4$ -space divided by $\mathbb{Z}/(4)$ with weights*

$\text{wt}(y_1, y_2, y_3, y_4) = (1, 3, 3, 2)$, and $K_Y = f^*K_X + 3E$. Moreover there exists such an example.

It is anticipated that the number of divisorial contractions over P decreases as the singularity of P becomes worse on account of restriction to choice of coordinates at P . Nevertheless we might not expect that it makes the perfect explicit study of them simple because the defining equation of X at P becomes much more complicated. By this reason, instead of complete description, we restrict the possibility of divisorial contractions in the remaining cD_n and cE_n cases by giving an upper bound of discrepancies (2.9), equipped with some typical examples.

Theorem 1.3. *Assume that P is cD_n or cE_n , and let $K_Y = f^*K_X + aE$. Then $a \leq 4$.*

We now have sufficient tools toward classification of divisorial contractions even in cD_n and cE_n cases once $P \in X$ is given explicitly. Basically what we should do is to compare discrepancies as in [Km] or in (7.1), with enormous information on multiplicities along E of surfaces on X with special directions provided by the singular Riemann-Roch technique and the general elephant theorem, occasionally applying Shokurov's connectedness lemma as in [Co, Theorem 3.10], [CM, Theorem 3.6].

This paper is constructed as follows. In Section 2 we state our theorems precisely. Section 3 is devoted to preparing basic numerical techniques. After recalling results for divisorial contractions to compound Du Val points in [Ka1], we introduce our fundamental setup toward analysis of $C \subset Y$ following [Mo2]. Quite delicate local investigation of $C \subset Y$ is presented in Section 4. Using it in Section 5 we prove our main theorem, the existence of Du Val sections. In Section 6 we restrict possible divisorial contractions focusing on the types of singularities on Du Val sections. It provides an upper bound of discrepancies in cD_n and cE_n cases. Finally in Section 7 we give explicit description of divisorial contractions in cA_n case.

I would like to thank Professor Yujiro Kawamata for his stimulating encouragement. I am grateful to Professor Alessio Corti and Professor Miles Reid for motivating me in this subject. I have written this paper during my visit to the University of Cambridge, where I receive warm hospitality of Professor Corti. Financial support has been provided by the Japan Society for the Promotion of Science.

2 Preliminaries and statements

We work over the complex number field $\mathbb{C}$. We fundamentally do in the analytic category but sometimes enter the algebraic category through algebraisation theorems of M. Artin [Ar1][Ar2]. First we define a divisorial contraction in a general sense.

Definition 2.1. Let $f: Y \rightarrow X$ be a morphism with connected fibres between normal varieties with at worst terminal singularities. We call f a *divisorial contraction* if the exceptional locus of f is a prime divisor and $-K_Y$ is f -ample.

Remark 2.1.1. Of course this definition applies to divisorial contractions emerging in the usual minimal model program [KMM].

We recall the classification of three dimensional terminal singularities.

Definition 2.2. Let $P \in X$ be a germ of a three dimensional variety. We call P a *cDV (compound Du Val) point* if a general hyperplane section has at worst a Du Val singularity at P . The singularity P is said to be cA_n , cD_n , cE_n (*compound A_n, D_n, E_n*) according to the type of the Du Val singularity on a general hyperplane section.

Remark 2.2.1. We say that a smooth point on a surface or a threefold is A_0 , cA_0 respectively for convenience.

Theorem 2.3 ([Re1, Theorem1.1]). *Let $P \in X$ be a germ of a three dimensional variety. Then P is a Gorenstein terminal singularity if and only if P is an isolated cDV point.*

2.4. Consider a non-Gorenstein terminal singularity $P \in X$. Let r be the local Gorenstein index of $P \in X$, that is, the smallest positive integer such that rK_X is Cartier at P . Take the index one cover $\pi: (X^\sharp \ni P^\sharp) \rightarrow (X \ni P)$, which is a cyclic μ_r -cover. Fix a character generating $\text{Hom}(\mu_r, \mathbb{C}^\times) = \mathbb{Z}/(r)$ and define the weight modulo r for any semi-invariant function on $X^\sharp$ with respect to this character.

Theorem 2.5 ([Mo1]). *There exists a μ_r -equivariant identification*

$$P^\sharp \in X^\sharp \cong o \in (\phi = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4\text{-space},$$

where x_1, x_2, x_3, x_4 and ϕ are μ_r -semi-invariant and $\phi = x_4$ if $P^\sharp \in X^\sharp$ is a smooth point. The weights of x_1, x_2, x_3, x_4 and ϕ satisfy one of the following.

- (i) $\text{wt}(x_1, x_2, x_3, x_4; \phi) = (1, -1, b, 0; 0)$, where b is coprime to r .
- (ii) $r = 4$ and $\text{wt}(x_1, x_2, x_3, x_4; \phi) = (1, 3, 3, 2; 2)$.

Remark 2.5.1. [Mo1] gives more precise description of ϕ and weights, and the classification is completed by [KS, Theorem 6.4].

Remark 2.5.2. Any three dimensional non-Gorenstein terminal singularity $P \in X$ has a small deformation to a basket of terminal quotient singularities P_i . P_i are called fictitious singularities in the sense of M. Reid [Re3]. If P is of type (i) in (2.5) then any local index at P_i equals that at P . If P is of type (ii) then the local index at one of P_i equals 2 and that at each of the rest equals 4.

Now it is the time when we state theorems. To begin with we give a numerical classification of f which is a slightly revised version of [Ka1, Theorem 4.5] quoted in (3.2), due to (3.8).

Theorem 2.6. *Let $f: (Y \supset E) \rightarrow (X \ni P)$ be a germ of a three dimensional divisorial contraction whose exceptional divisor E contracts to a cDV point P . Let $K_Y = f^*K_X + aE$, let $I = \{Q : \text{type } \frac{1}{r_Q}(1, -1, \overline{av_Q})\}$ ($v_Q \leq r_Q/2$) be the set of fictitious singularities from non-Gorenstein singularities on Y , let $J = \{(r_Q, v_Q)\}_{Q \in I}$, and let $1 + d(-1) = \dim \mathcal{O}_X / f_* \mathcal{O}_Y(-2E)$. Precise notation is provided in (3.1). Then f is exactly one of the following types.*

type	$1 + d(-1)$	J	a
O	≥ 2		1
I	1	$\{(7, 3)\}$ or $\{(3, 1), (5, 2)\}$	2
IIa	2	$\{(r, 2)\}$	2 or 4
IIb	2	$\{(r_1, 1), (r_2, 1)\}$	$(r_1 + r_2)/r_1 r_2 E^3 \geq 2$
III	3	$\{(r, 1)\}$	$(1 + r)/r E^3 \geq 2$
IV	4	$\emptyset$	2

Remark 2.6.1. For convenience we divide type IIb into two types according to the number of non-Gorenstein points on Y . We say that f is of type $IIb^\vee$ or $IIb^{\vee\vee}$ if f is of type II with one, two non-Gorenstein points on Y respectively.

Our main theorem is on general elements in the anticanonical system $|-K_Y|$ of Y , dubbed general elephants by M. Reid [Re3].

Theorem 2.7. *Let $f: (Y \supset E) \rightarrow (X \ni P)$ be a germ of a divisorial contraction in (2.6). Then a general element in $|-K_Y|$ has at worst Du Val singularities.*

Let S be a surface on Y defined by a general element in $|-K_Y|$ and let S_X be its strict transform on X . S and S_X have at worst Du Val singularities and the induced morphism $S \rightarrow S_X$ factors the minimal resolution of S_X by (2.7). We obtain the main theorem together with information on the partial resolution $S \rightarrow S_X$. (2.7) and (2.8) are induced by (3.8), (5.4), (5.5), (6.6), (6.9) with (3.2.1) and (6.9.1).

Theorem 2.8. (i) Assume that f is of type O or I. Then S_X defines a general element in $|-K_X|$. Moreover if f is of type I then P is cE_7 or cE_8 .

(ii) Assume that f is of type II, III or IV. Then possible types of Du Val singularities on S and S_X are as follows.

type of f	type of S_X	type of S
IIa	D_r or D_{r+1}	A_{r-1}
$IIb^\vee, J = \{(r, 1), (r, 1)\}$	D_{2r} or D_{2r+1}	A_{2r-1}
$IIb^\vee, J = \{(3, 1), (3, 1)\}$	E_7	E_6
$IIb^\vee, J = \{(2, 1), (4, 1)\}$	E_6	D_5
$IIb^{\vee\vee}$	$A_{r_1+r_2-1}$	A_{r_1-1} and A_{r_2-1}
III	A_r	A_{r-1}
IV	smooth	smooth

The type of a Du Val singularity $Q \in S$ for any point Q is that of a general Du Val section of a germ $Q \in Y$. Furthermore the exceptional locus of the partial resolution $S \rightarrow S_X$ is irreducible unless the type of S_X is that after the word “or” in the above table.

Remark 2.8.1. We have examples of type I with $J = \{(3, 1), (5, 2)\}$ in (6.7) but I do not know whether type I with $J = \{(7, 3)\}$ happens or not. If f is of type I with $J = \{(7, 3)\}$ then P is cE_7 (6.6.1).

We bound the discrepancy a in the case where P is cD_n or cE_n as a corollary.

Corollary 2.9. Assume that P is cD_n or cE_n . Then f is of type O, I, IIa or $IIb^\vee$ in (2.6) and the discrepancy $a \leq 4$.

Remark 2.9.1. We have examples of type O, I, $IIb^\vee$, or with $a = 1, 2, 3$ in (3.3), (6.7), (6.10) and (6.11). However I do not know whether type IIa happens or not. If $a = 4$ in cD_n or cE_n case then $r = 5$ and P has to be cD_4 , cD_5 or cD_6 (4.5(iii)).

Divisorial contractions to smooth or cA_1 points have been completely classified in [Ka1] and [Ka2].

Theorem 2.10 ([Ka1]). Assume that P is smooth. Then f is a weighted blowup. More precisely, we can take local coordinates x_1, x_2, x_3 at P and coprime positive integers s and t , such that f is the weighted blowup of X with weights $\text{wt}(x_1, x_2, x_3) = (1, s, t)$. Moreover any such f is a divisorial contraction.

Remark 2.10.1. We may assume that $s \leq t$. The type of f is $\text{Ib}^{\vee\vee}$, III, IV if $s > 1$, $s = 1$ and $t > 1$, $t = 1$ respectively.

Theorem 2.11 ([Ka2]). Assume that P is cA_1 . Then f is a weighted blowup. More precisely, under a suitable identification $P \in X \cong o \in (x_1x_2 + x_3^2 + x_4^N = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4$ -space, f is the weighted blowup with one of the following weights.

- (i) $\text{wt}(x_1, x_2, x_3, x_4) = (s, 2t - s, t, 1)$, where s, t are coprime positive integers such that $s \leq t \leq N/2$.
- (ii) $N = 3$ and $\text{wt}(x_1, x_2, x_3, x_4) = (1, 5, 3, 2)$.

Moreover any such f is a divisorial contraction.

Remark 2.11.1. In (i) the type of f is $\text{Ib}^{\vee\vee}$, III, O if $s > 1$, $s = 1$ and $t > 1$, $t = 1$ respectively. In (ii) the type of f is IIa.

Remark 2.11.2. A. Corti has obtained the result in the case where P is an ordinary double point earlier [Co, Theorem 3.10].

The case where P is cA_n ($n \geq 2$) is done in Section 7 in this paper.

Theorem 2.12. Assume that P is cA_n ($n \geq 2$). Then one of the following holds.

- (i) Under a suitable identification $P \in X \cong o \in (x_1x_2 + g(x_3, x_4) = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4$ -space, f is the weighted blowup with weights $\text{wt}(x_1, x_2, x_3, x_4) = (r_1, r_2, a, 1)$, where a divides $r_1 + r_2$ and is coprime to r_1 and r_2 , the weighted order of g with weights $\text{wt}(x_3, x_4) = (a, 1)$ is $r_1 + r_2$, and the coefficient of $x_3^{(r_1+r_2)/a}$ in g is not zero. Moreover any such f is a divisorial contraction.
- (ii) P is a cA_2 point isomorphic to $o \in (x_1x_2 + x_3^3 + g_{\geq 4}(x_3, x_4) = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4$ -space, where the total order of $g_{\geq 4}$ with respect to x_3, x_4 is greater than or equal to 4, Y has exactly one non-Gorenstein point Q , which is isomorphic to $o \in (y_1^2 + y_2^2 + y_3^2 + y_4^3 = 0)$ in the quotient space of $\mathbb{C}^4 = y_1y_2y_3y_4$ -space divided by $\mathbb{Z}/(4)$ with weights $\text{wt}(y_1, y_2, y_3, y_4) = (1, 3, 3, 2)$, and $K_Y = f^*K_X + 3E$. Moreover there exists such an example (7.10).

Remark 2.12.1. In (i) we may assume that $r_1 \leq r_2$. The type of f is $\text{Ib}^{\vee\vee}$, III, O if $r_1 > 1$ and $a > 1$, $r_1 = 1$ and $a > 1$, $a = 1$ respectively. In (ii) the type of f is $\text{Ib}^{\vee}$ with $J = \{(2, 1), (4, 1)\}$.

Remark 2.12.2. A. Corti and M. Mella have obtained the result in the case where P is a cA_2 point isomorphic to $o \in (x_1x_2 + x_3^3 + x_4^3 = 0) \subset \mathbb{C}^4$ earlier [CM, Theorem 3.6].

3 Basic numerical results

3.1. Let $f: (Y \supset E) \rightarrow (X \ni P)$ be a germ of a three dimensional divisorial contraction whose exceptional divisor E contracts to a cDV point P . We remark that f can be always extended to a morphism between projective varieties [Ar1][Ar2]. Let $K_Y = f^*K_X + aE$ and let r be the global Gorenstein index of Y .

Lemma 3.1.1 ([Ka1, Lemma 4.3]). *a and r are coprime.*

We take an integer e such that $ae \equiv 1$ modulo r . Let $I = \{Q : \text{type } \frac{1}{r_Q}(1, -1, b_Q)\}$ be the set of fictitious singularities from non-Gorenstein singularities on Y (2.5.2). Then $(\mathcal{O}_{Y_Q}(E_Q))_Q \cong (\mathcal{O}_{Y_Q}(eK_{Y_Q}))_Q$, where (Y_Q, E_Q) is the deformed pair at Q from (Y, E) . We note that b_Q is coprime to r_Q and that e is also coprime to r_Q because $ae \equiv 1$ modulo r . Hence $v_Q = \overline{eb_Q}$ is coprime to r_Q . Here $\overline{}$ denotes the smallest residue modulo r_Q , that is, $\overline{j} = j - \lfloor \frac{j}{r_Q} \rfloor r_Q$, where $\lfloor \rfloor$ denotes the round down, that is, $\lfloor j \rfloor = \max\{k \in \mathbb{Z} \mid k \leq j\}$. Replacing b_Q with $r_Q - b_Q$ if necessary, we may assume that $v_Q \leq r_Q/2$. With this description, $r = 1$ if I is empty, and otherwise r is the lowest common multiple of $\{r_Q\}_{Q \in I}$. We set $J = \{(r_Q, v_Q)\}_{Q \in I}$. We moreover define $d(i) = \dim f_*\mathcal{O}_Y(iE)/f_*\mathcal{O}_Y((i-1)E)$. We note that $d(i) = 0$ ($i \geq 1$) and $d(0) = 1$.

Remark 3.1.2. $a < \max\{r_Q\}$ unless P is a smooth point by analogue to the proof of [Ka2, Lemma 6.10], because there exists a valuation with centre P whose discrepancy with respect to K_X is 1 [Ma].

The next theorem gives a rough numerical classification of f .

Theorem 3.2 ([Ka1, Theorem 4.5]). *f is exactly of one of the following types.*

<i>type</i>	$1 + d(-1)$	J	a
<i>O</i>			1
<i>I</i>	1	$\{(7, 3)\}$ or $\{(3, 1), (5, 2)\}$	2
<i>IIa</i>	2	$\{(r, 2)\}$	$4/rE^3 = 2$ or 4
<i>IIb</i>	2	$\{(r_1, 1), (r_2, 1)\}$ ($r_1 \leq r_2$)	$(r_1 + r_2)/r_1r_2E^3 \geq 2$
<i>III</i>	3	$\{(r, 1)\}$	$(1 + r)/rE^3 \geq 2$
<i>IV</i>	4	$\emptyset$	2

Remark 3.2.1. f is of type IV if and only if f is the usual blowup along a smooth point P .

Remark 3.2.2. Type IIb is divided into types $\text{IIb}^\vee$ and $\text{IIb}^{\vee\vee}$ (2.6.1). By (2.5.2) if f is of type $\text{IIb}^\vee$ then $J = \{(r, 1), (r, 1)\}$ or $\{(2, 1), (4, 1)\}$, and $a = 2, 3$ respectively. In this case from [Mo1] the unique non-Gorenstein point $Q \in Y$ can be described as one of the following equations $\phi = 0$ using semi-invariant local coordinates y_1, y_2, y_3, y_4 of the index one cover $Q^\# \in Y^\#$ with weights $w = \text{wt}(y_1, y_2, y_3, y_4)$.

(i) $J = \{(r, 1), (r, 1)\}$.

(a) $\phi = y_1 y_2 + g(y_3^r, y_4)$ and $w = (1, -1, 2, 0)$. The order of $g(0, y_4)$ is 2. The type of a general Du Val section of a germ $Q \in Y$ is A_{2r-1} .

(b) $r = 3$, $\phi = y_4^2 + \phi_3(y_1, y_2, y_3) + \phi_{\geq 4}(y_1, y_2, y_3)$ and $w = (1, -1, 2, 0)$. ϕ_3 is $y_1^3 + y_2^3 + y_3^3$, $y_1^3 + y_2 y_3^2$ or $y_1^3 + y_2^3$, and $\phi_{\geq 4}$ is of order ≥ 4 . The type of a general Du Val section of a germ $Q \in Y$ is E_6 .

(ii) $J = \{(2, 1), (4, 1)\}$. $\phi = y_1^2 + y_2^2 + g(y_3^2, y_4)$ and $w = (1, 3, 3, 2)$. The order of $g(0, y_4)$ is 3. The type of a general Du Val section of a germ $Q \in Y$ is D_5 .

Remark 3.2.3. By the proof of this theorem in [Ka1], we can restrict possible values of J even if f is of type O . In fact J is one of

$\{(7, 3)\}$, $\{(8, 3)\}$,
 $\{(2, 1), (5, 2)\}$, $\{(3, 1), (5, 2)\}$, $\{(4, 1), (5, 2)\}$, $\{(2, 1), (7, 2)\}$,
 $\{(2, 1), (2, 1), (r_3, 1)\}$, $\{(2, 1), (3, 1), (3, 1)\}$,
 $\{(2, 1), (3, 1), (4, 1)\}$, $\{(2, 1), (3, 1), (5, 1)\}$,
 $\{(r, 2)\}$, $\{(r_1, 1), (r_2, 1)\}$, $\{(r, 1)\}$ and $\emptyset$.

In particular the number of fictitious singularities is at most three.

Example 3.3. The weighted blowup of the cD_4 singularity $o \in (x_1^2 + x_2^3 + x_3^3 + x_4^6 = 0) \subset \mathbb{C}^4$ with weights $\text{wt}(x_1, x_2, x_3, x_4) = (3, 2, 2, 1)$ in [Ka1, Example 4.6] is an example whose J is $\{(2, 1), (2, 1), (2, 1)\}$.

3.4. (3.2) is obtained from the singular Riemann-Roch formula [Re3, Theorem 10.2] and a relative vanishing theorem [KMM, Theorem 1-2-5] with the exact sequences

$$(3.4.1) \quad 0 \rightarrow \mathcal{O}_Y((i-1)E) \rightarrow \mathcal{O}_Y(iE) \rightarrow \mathcal{Q}_i \rightarrow 0.$$

$\mathcal{Q}_i$ are $\mathcal{S}_2$ by [KM2, Proposition 5.26], and are reflexive because they are locally free on the restriction $E^o \subseteq E$ of the Gorenstein locus of Y . Thus, since E is Cohen-Macaulay, $\mathcal{Q}_i$ are concretely given by $i_*(\mathcal{O}_Y(iE) \otimes \mathcal{O}_{E|E^o})$, where i is the induced map $E^o \hookrightarrow E$.

The singular Riemann-Roch implies

$$\begin{aligned}
(3.4.2) \quad \chi(\mathcal{Q}_i) &= \frac{1}{12} \{2(3i^2 - 3i + 1) - 3(2i - 1)a + a^2\} E^3 \\
&\quad + \frac{1}{12} E \cdot c_2(Y) + A_i - A_{i-1}, \\
A_i &= \sum_{Q \in I} \left(-\overline{ie} \frac{r_Q^2 - 1}{12r_Q} + \sum_{j=1}^{\overline{ie}-1} \frac{j\overline{b_Q}(r_Q - j\overline{b_Q})}{2r_Q} \right).
\end{aligned}$$

On the other hand a relative vanishing theorem implies

$$(3.4.3) \quad R^j f_* \mathcal{O}_Y(iE) = 0 \quad (i \leq a, j \geq 1).$$

We summarise the formulae in [Ka1] obtained from the above.

Proposition 3.5. (i) $rE^3 \in \mathbb{Z}_{>0}$.

(ii) $d(i) = \chi(\mathcal{Q}_i)$ ($i \leq a$).

(iii) If f is of type IIb or III and we set $r_1 = 1$, $r_2 = r$ when f is of type III, then

$$d(-i) = 1 + \left\lfloor \frac{i}{r_1} \right\rfloor \quad (0 \leq i < \min\{r_2, a\}).$$

(iv) Set $B_i = \sum_{Q \in I} \frac{\overline{iv_Q}(r_Q - \overline{iv_Q})}{2r_Q}$. Then

$$\chi(\mathcal{Q}_{-i}) - \chi(\mathcal{Q}_{i+1}) = \left(i + \frac{1}{2}\right) aE^3 + B_{i+1} - B_i.$$

Remark 3.5.1. The values of $d(-i) = \chi(\mathcal{Q}_{-i})$ below are used later.

(i) (type IIa) By direct calculation using (3.4.2), we have $d(-1) = 1$, $d(-2) \geq 2$ if $a = 2$ or $(a, r) = (4, 5)$.

(ii) (types IIb and III) $d(-i) = 1 + \left\lfloor \frac{i}{r_1} \right\rfloor$ ($1 \leq i < \min\{r_2, a\}$) (3.5(iii)), and $d(-a) = 2 + \left\lfloor \frac{a}{r_1} \right\rfloor$ if $a < r_2$ by (3.6) and the proofs of [Ka1, Proposition 4.4.3] and the latter part of [Ka1, Theorem 4.5].

Lemma 3.6. $\mathcal{Q}_{a+1} = \omega_E$ and $h^0(\omega_E) = h^1(\omega_E) = 0$, $h^2(\omega_E) = 1$.

Proof. Let E° on E be the restriction of the Gorenstein locus on Y and let $i: E^\circ \hookrightarrow E$ be the induced map. $\omega_E = i_*\omega_{E^\circ}$ since ω_E is S_2 . $\mathcal{Q}_{a+1} = \omega_E$ comes from $\mathcal{Q}_{a+1}|_{E^\circ} = \omega_{E^\circ}$.

$H^i(\omega_E)$ is the dual of $\text{Ext}_E^{2-i}(\omega_E, \omega_E)$. Therefore $h^0(\omega_E) = h^1(\omega_E) = 0$ since E is Cohen-Macaulay. $h^2(\omega_E) = \dim \text{Hom}(\omega_E, \omega_E)$. Since ω_E is S_2 , the map $\text{Hom}(\omega_E, \omega_E) \rightarrow \text{Hom}(\omega_{E^\circ}, \omega_{E^\circ})$ is injective. On the other hand $\dim \text{Hom}(\omega_{E^\circ}, \omega_{E^\circ}) = h^0(\mathcal{O}_{E^\circ}) = 1$. Because $\text{Hom}(\omega_E, \omega_E) \neq 0$, we obtain $h^2(\omega_E) = 1$. $\square$

3.7. Let $H_X \ni P$ be a general hyperplane section on X and let H be its strict transform on Y . Write $f^*H_X = H + bE$. b is the largest integer satisfying that $f_*\mathcal{O}_Y(-bE) = \mathfrak{m}_P$. Let Y° be the Gorenstein locus of Y . By the adjunction formula we have

$$\omega_H|_{Y^\circ \cap H} = f^*\omega_{H_X} \otimes \mathcal{O}_Y((a-b)E)|_{Y^\circ \cap H}.$$

$a \geq b$ because H_X is canonical. Moreover if $a = b$ then H is normal and gives a Du Val section in the anticanonical system of Y . Combining it with (3.2) we have the following.

Theorem 3.8. *If f is of type O or I , then the strict transform H of a general hyperplane section H_X on X gives a Du Val section in the anticanonical system on Y .*

3.9. In the remainder of this section we investigate the scheme $H \cap E$ of dimension 1. We use the commutative diagrams below repeatedly.

$$\begin{array}{ccccccc}
& \mathcal{O}_Y(iE) \otimes \mathcal{O}_Y(-E) & \rightarrow & \mathcal{O}_Y(iE) & \rightarrow & \mathcal{O}_Y(iE) \otimes \mathcal{O}_E & \rightarrow 0 \\
& \downarrow & & \parallel & & \downarrow & \\
0 \rightarrow & \mathcal{O}_Y((i-1)E) & \rightarrow & \mathcal{O}_Y(iE) & \rightarrow & \mathcal{Q}_i & \rightarrow 0, \\
& & & & & \downarrow & \\
& & & & & 0 & \\
& \uparrow & & & & & \\
& \mathcal{O}_Y(iE) \otimes \mathcal{I}_{H \cap E \subset E} & \rightarrow & \mathcal{O}_Y(iE) \otimes \mathcal{O}_E & \rightarrow & \mathcal{O}_Y(iE) \otimes \mathcal{O}_{H \cap E} & \rightarrow 0 \\
& \uparrow & & \parallel & & \parallel & \\
\mathcal{O}_Y(iE) \otimes \mathcal{O}_Y(-H) \otimes \mathcal{O}_E & \rightarrow & \mathcal{O}_Y(iE) \otimes \mathcal{O}_E & \rightarrow & \mathcal{O}_Y(iE) \otimes \mathcal{O}_{H \cap E} & \rightarrow 0 \\
& \downarrow & & \downarrow & & \downarrow & \\
0 \rightarrow & \mathcal{Q}_{i+b} & \rightarrow & \mathcal{Q}_i & \rightarrow & \mathcal{R}_i & \rightarrow 0, \\
& & & \downarrow & & \downarrow & \\
& & & 0 & & 0 &
\end{array}$$

where b is that in (3.7).

Chasing the diagrams (3.9) with (3.4.3) and (3.6), we have the following.

Lemma 3.10. (i) $h^j(\mathcal{O}_Y(iE) \otimes \mathcal{O}_E) = 0$ ($i \leq a, j \geq 1$).
 $h^1(\mathcal{O}_Y((a+1)E) \otimes \mathcal{O}_E) = 0$, $h^2(\mathcal{O}_Y((a+1)E) \otimes \mathcal{O}_E) = 1$.
(ii) If f is not of type I, then
 $h^1(\mathcal{O}_Y(iE) \otimes \mathcal{O}_{H \cap E}) = 0$ ($i \leq a-1$).
 $h^1(\mathcal{O}_Y(aE) \otimes \mathcal{O}_{H \cap E}) = 1$.

Remark 3.10.1. $h^0(\mathcal{O}_{H \cap E}) = 1$ and $h^1(\mathcal{O}_{H \cap E}) = 0$ in any case, even if f is of type I (3.8). Especially $H \cap E$ has no embedded points and $(H \cap E)_{\text{red}}$ is a union of $\mathbb{P}^1$.

3.11. The main ingredient we study is an irreducible reduced subscheme $C \cong \mathbb{P}^1 \subseteq H \cap E$. Here we introduce a normal form of $Q \in C \subset Y$ at each non-Gorenstein point Q of Y through which C passes, following [Mo2, Lemma 2.7].

Let r_Q be the local index of $Q \in Y$. Take the index one cover $\pi: (Y^\# \ni Q^\#) \rightarrow (Y \ni Q)$ and set $C^\# = (C \times_Y Y^\#)_{\text{red}}$. We have an identification (2.5)

$$Q^\# \in Y^\# \cong o \in (\phi = 0) \subset \mathbb{C}^4 = x_1 x_2 x_3 x_4\text{-space}.$$

Let s_Q be the number of irreducible components of $C^\#$ and $Q^\dagger \in C^\dagger$ be the normalisation of one of the irreducible components of $C^\#$. Let $t \in \mathcal{O}_{C, Q}$ and $t^{s_Q/r_Q} \in \mathcal{O}_{C^\dagger, Q^\dagger}$ be uniformising parameters of C and $C^\dagger$. Let a_i be the minimal number such that there exists a semi-invariant function with weight $\text{wt} x_i$ whose image in $\mathcal{O}_{C^\dagger, Q^\dagger}$ has order a_i/r_Q with respect to t . We note that

$$(3.11.1) \quad (a_1, a_2, a_3, a_4) \in s_Q \mathbb{Z}(\text{wt} x_1, \text{wt} x_2, \text{wt} x_3, \text{wt} x_4) \quad \text{in } (\mathbb{Z}/(r_Q))^4.$$

Then we can take an identification (2.5) such that $x_i|_{C^\dagger} = t^{a_i/r_Q}$ ($i = 1, 2, 3$), and $x_4|_{C^\dagger} = t^{a_4/r_Q}$ if $Q^\# \in Y^\#$ is singular.

3.12. We give how to compute the images of some natural maps of sheaves defined on a germ $Q \in C \subset Y$ using the data a_1, a_2, a_3, a_4 in (3.11). For $n \in \mathbb{Z}/(r_Q)$, we define $w_Q^C(n)$ as the smallest nonnegative integer such that $(w_Q^C(n), n) \in \mathbb{Z} \times \mathbb{Z}/(r_Q)$ is contained in the semigroup

$$(3.12.1) \quad \mathbb{Z}_{\geq 0}(a_1, \text{wt} x_1) + \mathbb{Z}_{\geq 0}(a_2, \text{wt} x_2) + \mathbb{Z}_{\geq 0}(a_3, \text{wt} x_3) + \mathbb{Z}_{\geq 0}(a_4, \text{wt} x_4),$$

or $\mathbb{Z}_{\geq 0}(a_1, \text{wt} x_1) + \mathbb{Z}_{\geq 0}(a_2, \text{wt} x_2) + \mathbb{Z}_{\geq 0}(a_3, \text{wt} x_3)$ if $Y^\#$ smooth.

$w_Q^C(0) = 0$, and $(r_Q, 0)$ is contained in the above semigroup because C is smooth. Take a reflexive sheaf $\mathcal{L}$ on Y which is isomorphic to the ideal sheaf defined by $x_1 = 0$ outside Q . We note that $\mathcal{L}^{[\otimes v_Q]} \cong \mathcal{O}_Y(E)$ and

$\mathcal{L}^{[\otimes b_Q]} \cong \mathcal{O}_Y(K_Y)$ on the germ at Q , where $\mathcal{L}^{[\otimes i]}$ denotes the double dual of $\mathcal{L}^{\otimes i}$. Then for any integers $j_1, \dots, j_k$ with $\sum_{1 \leq i \leq k} j_i = 0$, the image of the natural map

$$\mathcal{L}^{[\otimes j_1]} \otimes \dots \otimes \mathcal{L}^{[\otimes j_k]} \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$$

is $(\mathfrak{m}_{Q \subset C})^{\sum_{1 \leq i \leq k} w_Q^C(-j_i)/r_Q}$, where $\mathfrak{m}_{Q \subset C}$ is the ideal sheaf of Q in C .

3.13. We also need to treat sheaves $\mathcal{I}_C/\mathcal{I}_C^{(2)}$ as in [Mo2, Section 2]. For any $C \cong \mathbb{P}^1 \subset H \cap E$ its ideal sheaf as the subscheme in Y is denoted by $\mathcal{I}_C$, and its symbolic 2-power is denoted by $\mathcal{I}_C^{(2)}$. Let $C^{(2)} \subset Y$ be the closed subscheme defined by $\mathcal{I}_C^{(2)}$. $C^{(2)} \subseteq 2H \cap 2E$. $H^1(\mathcal{O}_{2H \cap 2E}) = 0$ unless f is of type O or I as in (3.10), and then $H^1(\mathcal{O}_{C^{(2)}}) = 0$. By the exact sequence

$$0 \rightarrow \mathcal{I}_C/\mathcal{I}_C^{(2)} \rightarrow \mathcal{O}_{C^{(2)}} \rightarrow \mathcal{O}_C \rightarrow 0,$$

we have

Lemma 3.13.1. $H^1(\mathcal{I}_C/\mathcal{I}_C^{(2)}) = 0$ unless f is of type O or I.

Consider the natural map

$$(3.13.2) \quad \bigwedge^2 \mathcal{I}_C/\mathcal{I}_C^{(2)} \otimes \mathcal{O}_C(K_C) \rightarrow [\mathcal{O}_Y(K_Y)]_C,$$

where $[\mathcal{L}]_C$ denotes the torsionfree part of $\mathcal{L} \otimes \mathcal{O}_C$. The length of the cokernel of this map is $2 + \deg[\mathcal{O}_Y(K_Y)]_C - \deg \bigwedge^2 \mathcal{I}_C/\mathcal{I}_C^{(2)}$. By (3.13.1) we obtain

Lemma 3.13.3. *The length of the cokernel of the map (3.13.2) is at most $4 + \deg[\mathcal{O}_Y(K_Y)]_C$ unless f is of type O or I.*

The next lemma is obtained by following [Mo2, Corollary 2.15] almost faithfully. We remark that $(2, 2) \in \mathbb{Z} \times \mathbb{Z}/(4)$ is contained in the semigroup (3.12.1) when Q is of exceptional type (ii) in (2.5) because C is smooth at Q .

Lemma 3.13.4 ([Mo2, Corollary 2.15]). *Let $Q \in C \subset Y$ be a singular point of Y . Then the map (3.13.2) is not surjective at Q .*

The next lemma demonstrates computation of the length of the cokernel of the map (3.13.2) in the simplest case.

Lemma 3.14. *Let $Q \in Y$ be a germ of a three dimensional terminal quotient singularity of index r obtained by the quotient of $\mathbb{C}^3 = x_1x_2x_3$ -space with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, b)$, and let $Q \in C \subset Y$ be a smooth curve with uniformising parameter t given by $(x_1, x_2, x_3)|_C = (t^{c/r}, t^{1-c/r}, 0)$ for some $0 < c < r$. Then the length of the cokernel of the map (3.13.2)*

$$\bigwedge^2 \mathcal{I}_C/\mathcal{I}_C^{(2)} \otimes \mathcal{O}_C(K_C) \rightarrow [\mathcal{O}_Y(K_Y)]_C$$

is $\min\{c, r - c\}$.

Proof. First we give an explicit description of this map using local coordinates. The map $\mathcal{O}_Y(K_Y) \otimes \mathcal{O}_C \rightarrow [\mathcal{O}_Y(K_Y)]_C \cong \mathcal{O}_C$ can be locally given, but not canonically, by

$$(3.14.1) \quad \mathcal{O}_Y(K_Y) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$$

$$f(x_1, x_2, x_3)dx_1 \wedge dx_2 \wedge dx_3 \mapsto \frac{f(t^{c/r}, t^{1-c/r}, 0)}{t^{w_Q^C(-b)/r}}.$$

$\mathcal{O}_C(K_C)$ is generated by $d(x_1x_2)$. Hence the length of the cokernel of our map equals that of the composition of

$$(3.14.2) \quad \mathcal{I}_C/\mathcal{I}_C^{(2)} \times \mathcal{I}_C/\mathcal{I}_C^{(2)} \rightarrow \mathcal{O}_Y(K_Y) \otimes \mathcal{O}_C$$

$$f_1 \times f_2 \mapsto df_1 \wedge df_2 \wedge d(x_1x_2)$$

and (3.14.1).

It is easy to see that $\mathcal{I}_C/\mathcal{I}_C^{(2)}$ is generated by functions of form $g_{-b}(x_1, x_2) \cdot x_3$ and $g_c(x_1, x_2) \cdot (x_1^{r-c} - x_2^c)$ where g_{-b} and g_c are semi-invariant functions of weights $-b$ and c respectively. Therefore we have only to consider images of $g_{-b}dx_3 \wedge d(g_c \cdot (x_1^{r-c} - x_2^c)) \wedge d(x_1x_2)$ by the map (3.14.1). Hence it is enough to consider $g_c = x_1^c$ and x_2^{r-c} . In these cases

$$g_{-b}dx_3 \wedge d(g_c \cdot (x_1^{r-c} - x_2^c)) \wedge d(x_1x_2)$$

$$= \begin{cases} g_{-b} \cdot rx_1^r dx_1 \wedge dx_2 \wedge dx_3 & (g_c = x_1^c) \\ g_{-b} \cdot rx_2^r dx_1 \wedge dx_2 \wedge dx_3 & (g_c = x_2^{r-c}). \end{cases}$$

Since we can choose g_{-b} so that $g_{-b}(t^{c/r}, t^{1-c/r}) = t^{w_Q^C(-b)/r}$, our length is the minimum of the orders of x_1^r and x_2^r with $(x_1, x_2) = (t^{c/r}, t^{1-c/r})$ with respect to t , which is $\min\{c, r-c\}$. $\square$

3.15. We need to compute the length in the more complicate case below. Let $Q \in Y$ be a germ of a three dimensional terminal quotient singularity of index r obtained by the quotient of $\mathbb{C}^3 = x_1x_2x_3$ -space with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, b)$, and let $Q \in C \subset Y$ be a smooth curve with uniformising parameter t given by $(x_1, x_2, x_3)|_C = (t^{a_1/r}, t^{a_2/r}, t^{a_3/r})$ for some a_1, a_2, a_3 . We can take an invariant monomial $g(x_1, x_2, x_3)$ such that $g(t^{a_1/r}, t^{a_2/r}, t^{a_3/r}) = t$.

$\mathcal{I}_C/\mathcal{I}_C^{(2)}$ is generated by functions of form $x_1^{s_1}x_2^{s_2}x_3^{s_3} - x_1^{t_1}x_2^{t_2}x_3^{t_3}$ with $(s_1, s_2, s_3) \neq (t_1, t_2, t_3)$ such that $s_1 - s_2 + bs_3 \equiv t_1 - t_2 + bt_3 \equiv 0$ modulo r and $a_1s_1 + a_2s_2 + a_3s_3 = a_1t_1 + a_2t_2 + a_3t_3$. Our length is the minimum of the orders of the images of $d(x_1^{s_1}x_2^{s_2}x_3^{s_3} - x_1^{t_1}x_2^{t_2}x_3^{t_3}) \wedge d(x_1^{\bar{s}_1}x_2^{\bar{s}_2}x_3^{\bar{s}_3} - x_1^{\bar{t}_1}x_2^{\bar{t}_2}x_3^{\bar{t}_3}) \wedge dg$ by the map

$$\mathcal{O}_Y(K_Y) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$$

$$f(x_1, x_2, x_3)dx_1 \wedge dx_2 \wedge dx_3 \mapsto \frac{f(t^{a_1/r}, t^{a_2/r}, t^{a_3/r})}{t^{w_Q^C(-b)/r}},$$

where $x_1^{s_1}x_2^{s_2}x_3^{s_3} - x_1^{t_1}x_2^{t_2}x_3^{t_3}$ and $x_1^{\bar{s}_1}x_2^{\bar{s}_2}x_3^{\bar{s}_3} - x_1^{\bar{t}_1}x_2^{\bar{t}_2}x_3^{\bar{t}_3}$ satisfy the conditions mentioned above. The order for fixed $x_1^{s_1}x_2^{s_2}x_3^{s_3} - x_1^{t_1}x_2^{t_2}x_3^{t_3}$ and $x_1^{\bar{s}_1}x_2^{\bar{s}_2}x_3^{\bar{s}_3} - x_1^{\bar{t}_1}x_2^{\bar{t}_2}x_3^{\bar{t}_3}$ is $+\infty$ or

$$\frac{1}{r} \cdot \{(a_1s_1 + a_2s_2 + a_3s_3) + (a_1\bar{s}_1 + a_2\bar{s}_2 + a_3\bar{s}_3) + r - (a_1 + a_2 + a_3 + w_Q^C(-b))\}.$$

In particular if we can choose $g = x_1x_2$ then it is $+\infty$ or

$$\frac{1}{r} \cdot \{(a_1s_1 + a_2s_2 + a_3s_3) + (a_1\bar{s}_1 + a_2\bar{s}_2 + a_3\bar{s}_3) - (a_3 + w_Q^C(-b))\}.$$

4 Local analysis at a non-Gorenstein point

4.1. In this section we analyse the structure of $C \cong \mathbb{P}^1 \subseteq H \cap E$ under the assumption that f is of type II or III focusing on the germ at a non-Gorenstein point. We can see the theorems in this section automatically when P is a smooth point because of the explicit description [Ka1], and hence we here impose an extra assumption, that is, P is a singular point. We have $(H \cdot C) \leq (H \cdot [H \cap E]) = E^3$. By (3.1.2), (3.2), (3.5(i)) and (3.10.1) the configuration of the 1-cycle $[H \cap E] = \sum [C_i]$ is as follows.

- (i) (type IIa) $a = 2$ or 4 and Y has one non-Gorenstein point Q , through which any C_i passes.
 - (a) $a = 2$ and $H \cap E \cong \mathbb{P}^1$ scheme theoretically.
 - (b) $a = 2$ and $[H \cap E] = [\mathbb{P}^1] + [\mathbb{P}^1]$ is reducible or nonreduced.
 - (c) $a = 4$ and $H \cap E \cong \mathbb{P}^1$ scheme theoretically.
- (ii) (type IIb $^\vee$) $J = \{(r, 1), (r, 1)\}$ or $\{(2, 1), (4, 1)\}$ and $a = 2, 3$ respectively. Y has one non-Gorenstein point Q . $H \cap E \cong \mathbb{P}^1$ scheme theoretically and passes through Q .
- (iii) (type IIb $^{\vee\vee}$) $a \leq (r_1 + r_2)/2$ and Y has two non-Gorenstein points Q_1 and Q_2 of indices r_1 and r_2 . Exactly one of the irreducible components of $(H \cap E)_{\text{red}}$ passes through both Q_1 and Q_2 , and the rest pass through Q_2 only.
- (iv) (type III) $a \leq (1+r)/2$ and Y has one non-Gorenstein point Q , through which any C_i passes.

4.2. Let $s_C(i)$ be the integer such that $[\mathcal{O}_Y(iE)]_C \cong \mathcal{O}_{\mathbb{P}^1}(s_C(i))$. The following are easy to see by (3.1.2), (3.10) and (4.3).

- (i) Assume that C comes from type IIa. Then

- (a) $s_C(1) = \cdots = s_C(a-1) = -1$.
 - (b) $s_C(-i) \geq -1$ ($i \geq 1$).
 - (c) $s_C(a) = -2$ in (4.1(ia), (ic)), and $s_C(a) = -2$ or -1 in (4.1(ib)).
- (ii) Assume that C comes from type $\text{IIb}^\vee$. Then
- (a) $s_C(1) = \cdots = s_C(a-1) = -1$. $s_C(a) = -2$.
 - (b) $s_C(-i) \geq -1$ ($i \geq 1$).
- (iii) Assume that C comes from type $\text{IIb}^{\vee\vee}$ with $Q_1, Q_2 \in C$. Then
- (a) $s_C(1) = \cdots = s_C(a-1) = -1$.
 - (b) $s_C(-i) = -1$ ($1 \leq i \leq a-1$, $r_1 \nmid i$).
 - (c) $s_C(-r_1 i) = 0$ or -1 ($1 \leq r_1 i \leq a-1$).
 - (d) $(s_C(a), s_C(-a)) = (-2, 0)$, $(-2, -1)$ or $(-1, -1)$.
- (iv) Assume that C comes from type III. Then
- (a) $s_C(1) = \cdots = s_C(a-1) = -1$.
 - (b) $s_C(-1) = \cdots = s_C(-(a-1)) = 0$.
 - (c) $(s_C(a), s_C(-a)) = (-2, 1)$, $(-2, 0)$ or $(-1, 0)$.

For instance we consider the case (iv). $s_C(i) < 0$ if $i \geq 1$ by the existence of the canonical map $\mathcal{O}_Y(iE)^{\otimes r} \otimes \mathcal{O}_C \rightarrow \mathcal{O}_Y(irE) \otimes \mathcal{O}_C$, and thus $s_C(1) = \cdots = s_C(a-1) = -1$, $s_C(a) = -2$ or -1 by (3.10). On the other hand (4.3) with $\mathcal{L} = \mathcal{O}_Y(E)$ implies that $s_C(-1) = 0$, and hence $s_C(-i) \geq 0$ if $i \geq 1$ by the existence of the canonical map $\mathcal{O}_Y(-E)^{\otimes i} \otimes \mathcal{O}_C \rightarrow \mathcal{O}_Y(-iE) \otimes \mathcal{O}_C$. Hence $s_C(-1) = \cdots = s_C(-(a-1)) = 0$ and $(s_C(a), s_C(-a)) = (-2, 1)$, $(-2, 0)$ or $(-1, 0)$ because the map $\mathcal{O}_Y(-iE) \otimes \mathcal{O}_Y(iE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ is not surjective when $1 \leq i \leq a < r$ (3.1.2).

Lemma 4.3. *Let $Q \in Y$ be a germ of a three dimensional terminal quotient singularity of index r obtained by the quotient of $\mathbb{C}^3 = x_1 x_2 x_3$ -space with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, b)$. Let $\mathcal{L}$ be a reflexive sheaf on Y which is isomorphic to the ideal sheaf defined by $x_1 = 0$ outside Q . Then the image of the natural map $\mathcal{L} \otimes \mathcal{L}^{[-1]} \rightarrow \mathcal{O}_Y$ is the ideal sheaf of Q in Y .*

Proof. It is enough to show that any invariant monomial $\neq 1$ of x_1, x_2, x_3 decomposes into two semi-invariant monomials with weights 1 and -1 , but it is trivial since b is coprime to r . $\square$

The following proposition stores more information on s_C .

Proposition 4.4. *Assume that f is not of type O or I , and take a curve $C \cong \mathbb{P}^1 \subseteq H \cap E$. Let i be a positive integer, and let T, T' be surfaces on Y defined by general elements in $\mathcal{O}_Y(-iE)$.*

- (i) *Assume that $\chi(\mathcal{Q}_i) \geq 2$. Then $(T \cap E)_{\text{red}}$ has an irreducible component which properly intersects $(T' \cap E)_{\text{red}}$.*
- (ii) *Assume that $\chi(\mathcal{Q}_j) = 1$ ($1 \leq j < i$) and $\chi(\mathcal{Q}_i) \geq 2$. If $C = H \cap E$ scheme theoretically, then $(T \cap E)_{\text{red}}$ properly intersects $(T' \cap E)_{\text{red}}$ and C . Moreover $s_C(-j) = -1$ ($1 \leq j < i$) and $s_C(-i) \geq 0$.*

Proof. (i) By (3.4.3) we have a surjective map $f_*\mathcal{O}_Y(-iE) \twoheadrightarrow H^0(\mathcal{Q}_{-i})$. Write E° as the restriction of the Gorenstein locus of Y to E . Take a resolution $\hat{E}$ of E and let $\hat{E}^\circ$ be the preimage of E° . We note that $H^0(\mathcal{Q}_{-i}|_{E^\circ}) = H^0(\mathcal{Q}_{-i})$ (3.4). Consider a linear system L on $\hat{E}$, not necessarily complete, whose restriction to $\hat{E}^\circ$ is $H^0(\mathcal{Q}_{-i}|_{E^\circ})$. $h^0(\mathcal{Q}_{-i}) = \chi(\mathcal{Q}_{-i}) \geq 2$ means that L moves, and its general member has an irreducible component which is not contained in the union of the support of the fixed locus of L and the preimage of the nonnormal locus of E . The element in $H^0(\mathcal{Q}_{-i})$ corresponding to this member gives a desired section T .

(ii) By (3.9) and (3.10(i)) we have surjective maps

$$f_*\mathcal{O}_Y(-jE) \twoheadrightarrow H^0(\mathcal{Q}_{-j}) \twoheadrightarrow H^0(\mathcal{R}_{-j}) \twoheadrightarrow H^0([\mathcal{O}_Y(-jE)]_C) \quad (1 \leq j \leq i).$$

We note that E is smooth at the generic point of C because $C = H \cap E$ scheme theoretically. $H^0(\mathcal{Q}_{-j}) = \chi(\mathcal{Q}_{-j}) = 1$ ($1 \leq j < i$) and a nonzero element in it corresponds to, at the generic point of C , the Cartier divisor defined by jH . Hence the map $H^0(\mathcal{Q}_{-j}) \twoheadrightarrow H^0([\mathcal{O}_Y(-jE)]_C)$ is a zero map, and we obtain $s_C(-j) = -1$ by the above map and (3.10(ii)).

On the other hand the L constructed from $H^0(\mathcal{Q}_{-i})$ as in the proof of (i) moves. Because $h^0(\mathcal{Q}_{-j}) = 1$ ($1 \leq j < i$) and E is smooth at the generic point of C , its general member does not have the pull-back of C as its component. Hence we have a desired section and we see that the map $H^0(\mathcal{Q}_{-i}) \twoheadrightarrow H^0([\mathcal{O}_Y(-iE)]_C)$ is a nonzero map, whence $s_C(-i) \geq 0$. $\square$

First we treat type IIa.

Theorem 4.5. *Assume that f is of type IIa. Let $Q \in C \cong \mathbb{P}^1 \subseteq H \cap E$ be the non-Gorenstein point with an irreducible component of $H \cap E$ (4.1(i)). Take a normal form of $Q \in C \subset Y$ (3.11).*

- (i) *The case (ia) in (4.1) does not happen.*
- (ii) *In the case (ib) in (4.1), we have $s_C(-2) = 0$ and $(a_1, a_2, a_3) = ((r+1)/2, (r-1)/2, 2)$. In particular we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, 4)$ of the index one cover $Q^\sharp \in Y^\sharp$ so that $(x_1, x_2, x_3)|_{C^\dagger} = (t^{(r+1)/2r}, 0, t^{2/r})$ ($r \equiv 1$ modulo 4) or $(0, t^{(r-1)/2r}, t^{2/r})$ ($r \equiv 3$ modulo 4).*

- (iii) In the case (ic) in (4.1), we have $r = 5$, $s_C(-4) = 0$ and $(a_1, a_2, a_3) = (3, 2, 4)$. In particular we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, 3)$ of the index one cover $Q^\# \in Y^\#$ so that $(x_1, x_2, x_3)|_{C^\dagger} = (t^{3/5}, t^{2/5}, 0)$.

Proof. (i). Compute the image of the map

$$\mathcal{O}_Y(E)^{\otimes r} \otimes \mathcal{O}_Y(-rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$$

by (3.12). It is $(\mathfrak{m}_{Q \subset C})^{w_Q^C(-2)}$. On the other hand this image is the same as that of

$$[\mathcal{O}_Y(E)]_C^{\otimes r} \otimes [\mathcal{O}_Y(-rE)]_C \rightarrow \mathcal{O}_C,$$

whence $rs_C(1) - r(E \cdot C) = -w_Q^C(-2)$. By $s_C(1) = -1$ (4.2(i)) and $(H \cdot C) = E^3 = 2/r$ (3.2) we have $w_Q^C(-2) = r - 2$. Hence we can write $(a_1, a_2, a_3) = (1 + rm_1, r - 1 + rm_2, 4 + rm_3)$ with $m_1, m_2, m_3 \in \mathbb{Z}_{\geq 0}$ by (3.11.1). We see $m_1 = 0$ easily since there exists an invariant monomial of x_1, x_2, x_3 whose restriction on C is t . Thus $(a_1, a_2, a_3) = (1, r - 1, 4)$ and especially $w_Q^C(-4) = r - 4$.

Consider another map $\mathcal{O}_Y(2E)^{\otimes r} \otimes \mathcal{O}_Y(-2rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$. Then we have $rs_C(2) - 2r(E \cdot C) = -w_Q^C(-4)$, whence $s_C(2) = -1$ by $w_Q^C(-4) = r - 4$ and $(H \cdot C) = 2/r$. But this contradicts (4.2(i)).

(ii) By the map $\mathcal{O}_Y(E)^{\otimes r} \otimes \mathcal{O}_Y(-rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we have $rs_C(1) - r(E \cdot C) = -w_Q^C(-2)$. Hence $w_Q^C(-2) = r - 1$ by (4.2(i)) and $(H \cdot C) = E^3/2 = 1/r$ (3.2), and we can write $(a_1, a_2, a_3) = ((r+1)/2 + rm_1, (r-1)/2 + rm_2, 2 + rm_3)$ with $m_1, m_2, m_3 \in \mathbb{Z}_{\geq 0}$ by (3.11.1). Because there exists an invariant monomial of x_1, x_2, x_3 whose restriction on C is t , we have one of the following.

- (i) $(r, 0) \in \mathbb{Z}_{>0}(a_1, 1) + \mathbb{Z}_{>0}(a_3, 4)$ or $\in \mathbb{Z}_{>0}(a_2, -1) + \mathbb{Z}_{>0}(a_3, 4)$ in $\mathbb{Z} \times \mathbb{Z}/(r)$.

- (ii) $r = a_1 + a_2$.

If (i) holds then $(a_1, a_2, a_3) = ((r+1)/2, (r-1)/2, 2)$ and $w_Q^C(4) = 4$. By the map $\mathcal{O}_Y(-2E)^{\otimes r} \otimes \mathcal{O}_Y(2rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we have $rs_C(-2) + 2r(E \cdot C) = -w_Q^C(4)$, whence $s_C(-2) = 0$. Now we assume (ii). Then $(a_1, a_2, a_3) = ((r+1)/2, (r-1)/2, 2 + rm_3)$ with some m_3 . We have nothing to do if $m_3 = 0$. Thus we also assume that $m_3 > 0$ and $r \geq 7$. We note that $m_3 \leq 2$ by $4a_1 = 2 + 2r$.

If $m_3 = 2$ then we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, 4)$ of $Q^\# \in Y^\#$ so that $(x_1, x_2, x_3)|_{C^\dagger} = (t^{(r+1)/2r}, t^{(r-1)/2r}, 0)$. We see $w_Q^C(-4) = 4w_Q^C(-1) = 2r - 2$ from $r \geq 7$. Considering the map $\mathcal{O}_Y(2E)^{\otimes r} \otimes \mathcal{O}_Y(-2rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we obtain $rs_C(2) - 2r(E \cdot C) = -w_Q^C(-4)$, whence $s_C(2) = -2$. By (3.13.3), the length

of the cokernel of the map (3.13.2) at Q is at most 2. Hence we have $(r-1)/2 \leq 2$ by (3.14), which contradicts $r \geq 7$.

Now it remains to exclude the case where $(a_1, a_2, a_3) = ((r+1)/2, (r-1)/2, r+2)$ with $r \geq 7$. But if $r = 7$ then $(a_1, a_2, a_3) = (4, 3, 9)$ and we may choose the coordinates so that $(x_1, x_2, x_3)|_{C^\dagger} = (t^{4/7}, t^{3/7}, 0)$. Then $s_C(2) = -2$ follows and contradicts (3.13.3) and (3.14). Therefore we may assume that $r \geq 9$. By the map $\mathcal{O}_Y(-3E)^{\otimes r} \otimes \mathcal{O}_Y(3rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we have $rs_C(-3) + 3r(E \cdot C) = -w_Q^C(6)$, whence $w_Q^C(6) = 3$ or $r+3$ since $s_C(-3) \geq -1$. From $(a_1, a_2, a_3) = ((r+1)/2, (r-1)/2, r+2)$ with $r \geq 9$, r has to be 9.

$(a_1, a_2, a_3) = (5, 4, 11)$ and $w_Q^C(-4) = 16$ when $r = 9$. By the map $\mathcal{O}_Y(2E)^{\otimes r} \otimes \mathcal{O}_Y(-18E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we obtain $9s_C(2) - 18(E \cdot C) = -w_Q^C(-4)$, whence $s_C(2) = -2$. We consider $x_1^{s_1}x_2^{s_2}x_3^{s_3} - x_1^{t_1}x_2^{t_2}x_3^{t_3}$ with $(s_1, s_2, s_3) \neq (t_1, t_2, t_3)$ such that $s_1 - s_2 + 4s_3 \equiv t_1 - t_2 + 4t_3 \equiv 0$ modulo 9 and $5s_1 + 4s_2 + 11s_3 = 5t_1 + 4t_2 + 11t_3$. We can easily see that $(5s_1 + 4s_2 + 11s_3)/9 \geq 3$ for any such $x_1^{s_1}x_2^{s_2}x_3^{s_3} - x_1^{t_1}x_2^{t_2}x_3^{t_3}$. Hence by (3.15), the length of the cokernel of the map (3.13.2) is greater than or equal to $3 + 3 - (a_3 + w_Q^C(-4))/9 = 3$, which contradicts $s_C(2) = -2$ and (3.13.3).

(iii) By the map $\mathcal{O}_Y(iE)^{\otimes r} \otimes \mathcal{O}_Y(-irE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ ($i = 1, 2, 3, 4$), we have $rs_C(i) - ir(E \cdot C) = -w_Q^C(-2i)$, whence $w_Q^C(-2i) = r - i$ ($i = 1, 2, 3$) and $w_Q^C(-8) = 2r - 4$ (4.2(ic)). We can write $(a_1, a_2, a_3) = ((r+1)/2 + rm_1, (r-1)/2 + rm_2, 4 + rm_3)$ with $m_1, m_2, m_3 \in \mathbb{Z}_{\geq 0}$ by (3.11.1). If $r = 5$ then $a_1 = w_Q^C(-4) = 3$ and $a_2 = w_Q^C(-6) = 2$ and we have the desired description. In this case $w_Q^C(8) = 4$, and by the map $\mathcal{O}_Y(-4E)^{\otimes r} \otimes \mathcal{O}_Y(4rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we have $rs_C(-4) + 4r(E \cdot C) = -w_Q^C(8)$, whence $s_C(-4) = 0$.

Now we derive a contradiction supposing that $r \geq 7$. Since $w_Q^C(-4) = r - 2$, we can decompose $r - 2$ into $(r+1)/2 + 4c$ or $(r-1)/2 + 4c$ for some positive integers c , and then $r \equiv 5, 3$ modulo 8 respectively. Since $w_Q^C(-6) = r - 3$, similarly we can decompose $r - 3$ into $(r+1)/2 + 4c$, $(r-1)/2 + 4c$ or $4c$ and then $r \equiv 7, 5$ modulo 8, $r \equiv 3$ modulo 4 respectively. Hence $r \equiv 3$ modulo 8 with $m_2 = m_3 = 0$, or $r \equiv 5$ modulo 8 with $m_1 = m_2 = m_3 = 0$. In particular $r \geq 11$ and $a_3 = 4$.

We consider $x_1^{s_1}x_2^{s_2}x_3^{s_3} - x_1^{t_1}x_2^{t_2}x_3^{t_3}$ with $(s_1, s_2, s_3) \neq (t_1, t_2, t_3)$ such that $s_1 - s_2 + 8s_3 \equiv t_1 - t_2 + 8t_3 \equiv 0$ modulo r and $a_1s_1 + a_2s_2 + a_3s_3 = a_1t_1 + a_2t_2 + a_3t_3$. We can easily see that $(a_1s_1 + a_2s_2 + a_3s_3)/r = 2$ only if each of $x_1^{s_1}x_2^{s_2}x_3^{s_3}$ and $x_1^{t_1}x_2^{t_2}x_3^{t_3}$ is $x_1^2x_2^2$, $x_1x_3^{(3r-1)/8}$ ($r \equiv 3$ modulo 8) or $x_2x_3^{(3r+1)/8}$ ($r \equiv 5$ modulo 8) and otherwise $(a_1s_1 + a_2s_2 + a_3s_3)/r \geq 3$. Hence by (3.15), the length of the cokernel of the map (3.13.2) is greater than or equal to $2 + 3 - (a_3 + w_Q^C(-8))/r = 3$, which contradicts $s_C(4) = -2$ (4.2(i)) and (3.13.3). $\square$

It is easy to see the local structure in type IIb $^\vee$.

Theorem 4.6. Assume that f is of type $\text{IIb}^\vee$. Let $Q \in C = H \cap E \cong \mathbb{P}^1$ be the non-Gorenstein point (4.1(ii)). Take a normal form of $Q \in C \subset Y$ (3.11).

- (i) If $J = \{(r, 1), (r, 1)\}$, then $a = 2$, $s_C(-2) = 0$ and $(a_1, a_2, a_3, a_4) = (r+1, r-1, 2, r)$. In particular we may choose semi-invariant local coordinates x_1, x_2, x_3, x_4 with weights $\text{wt}(x_1, x_2, x_3, x_4) = (1, -1, 2, 0)$ of the index one cover $Q^\sharp \in Y^\sharp$ so that $(x_1, x_2, x_3, x_4)|_{C^\dagger} = (0, 0, t^{2/r}, t)$.
- (ii) If $J = \{(2, 1), (4, 1)\}$, then $a = 3$, $s_C(-3) = 0$ and $(a_1, a_2, a_3, a_4) = (5, 3, 3, 2)$. In particular we may choose semi-invariant local coordinates x_1, x_2, x_3, x_4 with weights $\text{wt}(x_1, x_2, x_3, x_4) = (1, 3, 3, 2)$ of the index one cover $Q^\sharp \in Y^\sharp$ so that $(x_1, x_2, x_3, x_4)|_{C^\dagger} = (0, 0, t^{3/4}, t^{1/2})$.

Proof. (i) First applying (4.4(ii)) with $i = 2$ (3.5.1(ii)) we have $s_C(-1) = -1$ and $s_C(-2) \geq 0$. The map $\mathcal{O}_Y(-2E) \otimes \mathcal{O}_Y(E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_Y(-E) \otimes \mathcal{O}_C$ implies that $s_C(-2) + s_C(1) \leq s_C(-1)$. Hence $s_C(-2) = 0$. By the map $\mathcal{O}_Y(-E)^{\otimes r} \otimes \mathcal{O}_Y(rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we have $rs_C(-1) + r(E \cdot C) = -w_Q^C(1)$. By $s_C(-1) = -1$ and $E^3 = 1/r$ we have $w_Q^C(1) = 1 + r$. Hence we can write $(a_1, a_2, a_3, a_4) = (r+1+rm_1, r-1+rm_2, 2+rm_3, r)$ with $m_1, m_2, m_3 \in \mathbb{Z}_{\geq 0}$ by (3.11.1) and the property that $(r, 0)$ is contained in the semigroup (3.12.1). Consider another map $\mathcal{O}_Y(-2E)^{\otimes r} \otimes \mathcal{O}_Y(2rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$. Then we have $rs_C(-2) + 2r(E \cdot C) = -w_Q^C(2)$. Therefore $w_Q^C(2) = 2$ from $s_C(-2) = 0$, whence $(a_1, a_2, a_3, a_4) = (r+1, r-1, 2, r)$.

(ii) By (4.4(ii)) with large i (3.5.1(ii)) we have $s_C(-1) = -1$. By the map $\mathcal{O}_Y(-E)^{\otimes 4} \otimes \mathcal{O}_Y(4E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we have $4s_C(-1) + 4(E \cdot C) = -w_Q^C(1)$. By $s_C(-1) = -1$ and $E^3 = 1/4$ we obtain $w_Q^C(1) = 5$. Hence we can write $(a_1, a_2, a_3, a_4) = (5+4m_1, 3+4m_2, 3+4m_3, 2)$ with $m_1, m_2, m_3 \in \mathbb{Z}_{\geq 0}$. Consider another map $\mathcal{O}_Y(E)^{\otimes 4} \otimes \mathcal{O}_Y(-4E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$. Then we have $4s_C(1) - 4(E \cdot C) = -w_Q^C(-1)$. Thus $w_Q^C(-1) = 3$ (4.2(ii)), whence $(a_1, a_2, a_3, a_4) = (5, 3, 3, 2)$. $w_Q^C(3) = 3$. By the map $\mathcal{O}_Y(-3E)^{\otimes 4} \otimes \mathcal{O}_Y(12E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we have $4s_C(-3) + 12(E \cdot C) = -w_Q^C(3)$, whence $s_C(-3) = 0$. $\square$

4.7. It is hardest to treat types $\text{IIb}^{\vee\vee}$ and III.

- (i) If f is of type $\text{IIb}^{\vee\vee}$, then there exists the unique irreducible component $C \cong \mathbb{P}^1 \subseteq H \cap E$ through Q_1 and Q_2 . We do not consider any other components from now on. Take normal forms of $Q_i \in C \subset Y$ ($i = 1, 2$) using local coordinates x_{i1}, x_{i2}, x_{i3} and numbers a_{i1}, a_{i2}, a_{i3} respectively (3.11).
- (ii) If f is of type III, then we take any irreducible component $C \cong \mathbb{P}^1 \subseteq H \cap E$ and a normal form of $Q \in C \subset Y$ (3.11).

Proposition 4.8. *Assume that $s_C(-a) = 0$ in (4.7(i)) or $s_C(-a) = 1$ in (4.7(ii)). Then $C = H \cap E$ scheme theoretically, $s_C(a) = -2$, and for any non-Gorenstein point $Q \in Y$ with $Q \in C \cong \mathbb{P}^1 = H \cap E$ in (4.7) we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, a)$ of the index one cover $Q^\sharp \in Y^\sharp$ so that*

$$Q^\sharp \in C^\sharp \subset Y^\sharp \cong o \in (x_3\text{-axis}) \subset x_1x_2x_3\text{-space}.$$

Proof. By the map $\mathcal{O}_Y(-aE)^{\otimes r} \otimes \mathcal{O}_Y(arE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we have

$$rs_C(-a) + ar(E \cdot C) = \begin{cases} -rw_{Q_1}^C(a)/r_1 - rw_{Q_2}^C(a)/r_2 & (4.7(i)) \\ -w_Q^C(a) & (4.7(ii)). \end{cases}$$

Therefore by (3.2),

$$\frac{w_{Q_1}^C(a)}{r_1} + \frac{w_{Q_2}^C(a)}{r_2} = a(H \cdot C) \leq \frac{1}{r_1} + \frac{1}{r_2} \quad (4.7(i)),$$

$$\frac{w_Q^C(a)}{r} = a(H \cdot C) - 1 \leq \frac{1}{r} \quad (4.7(ii)).$$

Hence $C = H \cap E$ (3.10.1) and $w_Q^C(a) = 1$ for any $Q \in C$ in (4.7), and we have the desired description. $s_C(a) = -2$ follows from (3.10(ii)). $\square$

Proposition 4.9. *Assume that f is of type $I\text{Ib}^{\vee\vee}$ or III . Consider a non-Gorenstein point $Q \in Y$ with $Q \in C \cong \mathbb{P}^1 \subseteq H \cap E$ given in (4.7). Then one of the following holds.*

- (i) *One of $(a_1, 1)$, $(a_2, -1)$ is generated by the other and (a_3, a) in $\mathbb{Z} \times \mathbb{Z}/(r_Q)$, and we may choose semi-invariant local coordinates x_1, x_2, x_3 with weight $\text{wt}(x_1, x_2, x_3) = (1, -1, a)$ of the index one cover $Q^\sharp \in Y^\sharp$ so that $(x_1, x_2, x_3)|_{C^\dagger} = (0, t^{c_Q/r_Q}, t^{1-\overline{ac_Q}/r_Q})$ or $(t^{c_Q/r_Q}, 0, t^{\overline{ac_Q}/r_Q})$ for some $0 < c_Q < r_Q$.*
- (ii) *f is of type III , $a = 2$, $r \geq 5$, $s_Q(2) = -2$ and $(a_1, a_2, a_3) = (2, r-2, 4)$ or $(r-2, 2, 2r-4)$. In particular we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, 2)$ of the index one cover $Q^\sharp \in Y^\sharp$ so that $(x_1, x_2, x_3)|_{C^\dagger} = (t^{2/r}, t^{1-2/r}, 0)$ or $(t^{1-2/r}, t^{2/r}, 0)$. The value of $(H \cdot E)$ is $2/r$, $1 - 2/r$ respectively.*

Proof. Recall that there exists an invariant monomial of x_1, x_2, x_3 whose restriction on C is t . Thus one of the following holds.

- (i) one of a_1, a_2, a_3 is 1.
- (ii) $(r_Q, 0) \in \mathbb{Z}_{>0}(a_1, 1) + \mathbb{Z}_{>0}(a_3, a)$ or $\in \mathbb{Z}_{>0}(a_2, -1) + \mathbb{Z}_{>0}(a_3, a)$ in $\mathbb{Z} \times \mathbb{Z}/(r_Q)$.

(iii) $r_Q = a_1 + a_2$.

If (i) or (ii) happens, then one of $(a_1, 1)$ and $(a_2, -1)$ is generated by the other and (a_3, a) . Therefore we may assume that only (iii) holds from now on. By permutation of x_i we may moreover assume that $r_Q \geq 5$. Then it is easy to see that $w_Q^C(i) + w_Q^C(-i) \neq r_Q$ ($i = 2, 3$), whence the image of $[\mathcal{O}_Y(-iE)]_C \otimes [\mathcal{O}_Y(iE)]_C \rightarrow \mathcal{O}_C$ at Q is properly contained in $\mathfrak{m}_{Q \subset C}$ ($i = 2, 3$) (3.12). By (4.2) one of the following occurs.

- (i) $a = 2$, $s_C(2) = -2$, $s_C(-2) = -1$ in (4.7(i)) or $s_C(-2) = 0$ in (4.7(ii)).
- (ii) $Q = Q_2$, $a = 3$, $r_1 = 2$, $s_C(3) = -2$ and $s_C(-2) = s_C(-3) = -1$ in (4.7(i)).

Consider the case (i). $w_Q^C(1) + w_Q^C(-1) = r_Q$ and $w_Q^C(2) + w_Q^C(-2) = 2r_Q$, whence $w_Q^C(2) = 2w_Q^C(1)$ and $w_Q^C(-2) = 2w_Q^C(-1)$. We have $(a_1, a_2, a_3) = (c, r_Q - c, 2c)$ by putting $w_Q^C(1) = c$, and thus we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, 2)$ of the index one cover $Q^\# \in Y^\#$ so that $(x_1, x_2, x_3)|_{C^\dagger} = (t^{c/r_Q}, t^{1-c/r_Q}, 0)$.

By $s_C(2) = -2$, (3.13.3) and (3.13.4), the length of the map (3.13.2) at Q is at most 1 in (4.7(i)) and at most 2 in (4.7(ii)). Hence we have the result by (3.14) except the value of $(H \cdot C)$. This value can be calculated from $rs_C(-1) + r(E \cdot C) = -w_Q^C(1)$ obtained by the map $\mathcal{O}_Y(-E)^{\otimes r} \otimes \mathcal{O}_Y(rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$.

Consider the case (ii). $w_Q^C(3) + w_Q^C(-3) = 2r_Q$. By $s_C(3) = -2$ and (3.13.3) and (3.13.4), the length of the map (3.13.2) at Q must be 1. On the other hand, consider $x_1^{s_1} x_2^{s_2} x_3^{s_3} - x_1^{t_1} x_2^{t_2} x_3^{t_3}$ with $(s_1, s_2, s_3) \neq (t_1, t_2, t_3)$ such that $s_1 - s_2 + 3s_3 \equiv t_1 - t_2 + 3t_3 \equiv 0$ modulo r_Q and $a_1 s_1 + a_2 s_2 + a_3 s_3 = a_1 t_1 + a_2 t_2 + a_3 t_3$. We can easily see that $(a_1 s_1 + a_2 s_2 + a_3 s_3)/r_Q \geq 2$ for any such $x_1^{s_1} x_2^{s_2} x_3^{s_3} - x_1^{t_1} x_2^{t_2} x_3^{t_3}$ because $x_1 x_2$ is the only invariant monomial whose restriction to $C^\dagger$ is t . Hence by (3.15), the length of the cokernel of the map (3.13.2) at Q is greater than or equal to $2 + 2 - (a_3 + w_Q^C(-3))/r_Q = 2$, which is a contradiction. $\square$

We need more accurate description developing (4.9).

Lemma 4.10. *Consider the case in (4.9(i)).*

- (i) *If f is of type $Ib^{\vee\vee}$ (4.7(i)), then we may choose the coordinates so that the restriction to $C^\dagger$ at one of Q_1 and Q_2 is of form $(0, t^{c_i/r_i}, t^{1-\overline{ac_i}/r_i})$ and that at the other is of form $(t^{c_i/r_i}, 0, t^{\overline{ac_i}/r_i})$.*
- (ii) *If f is of type III (4.7(ii)), then we may choose the coordinates so that the restriction to $C^\dagger$ at Q is of form $(t^{c/r}, 0, t^{\overline{ac}/r})$.*

Proof. (i) If they both are of form $(0, t^{c_i/r_i}, t^{1-\overline{ac_i}/r_i})$, then the image of the map $\mathcal{O}_Y(-aE) \otimes \mathcal{O}_Y(E) \otimes \mathcal{O}_Y((a-1)E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ is $\mathfrak{m}_{\{Q_1, Q_2\} \subset C}$, whence

$s_C(-a) + s_C(1) + s_C(a-1) = -2$. Thus $s_C(-a) = 0$ (4.2(iii)), and it is done by (4.8). If they both are of form $(t^{c_i/r_i}, 0, t^{\overline{ac_i}/r_i})$, then the image of the map $\mathcal{O}_Y(-aE) \otimes \mathcal{O}_Y(-E) \otimes \mathcal{O}_Y((a+1)E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ is $\mathfrak{m}_{\{Q_1, Q_2\} \subset C}$, whence $s_C(-a) + s_C(-1) + s_C(a+1) = -2$. Thus $s_C(-a) = 0$ by $s_C(a+1) \leq -1$ and (4.2(iii)), and it is done by (4.8).

(ii) If the restriction is of form $(0, t^{c/r}, t^{1-\overline{ac}/r})$, then the image of the map $\mathcal{O}_Y(-aE) \otimes \mathcal{O}_Y(E) \otimes \mathcal{O}_Y((a-1)E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ is $\mathfrak{m}_{Q \subset C}$, whence $s_C(-a) + s_C(1) + s_C(a-1) = -1$. Thus $s_C(-a) = 1$ (4.2(iv)), and it is done by (4.8). $\square$

Proposition 4.11. (i) Assume that f is of type $Ib^{\vee\vee}$ with $s_C(-a) = -1$ (4.7(i)). Then $a < r_1 < r_2$, $a_{11} = w_{Q_1}^C(1) = 1$, $a_{22} = w_{Q_2}^C(-1) = 1$ and $(H \cdot C) = 1/r_1 - 1/r_2$. In particular for each i we may choose semi-invariant local coordinates x_{i1}, x_{i2}, x_{i3} with weights $\text{wt}(x_{i1}, x_{i2}, x_{i3}) = (1, -1, a)$ of the index one cover $Q_i^\sharp \in Y^\sharp$ so that

$$Q_i^\sharp \in C^\sharp \subset Y^\sharp \cong o \in (x_{ii}\text{-axis}) \subset x_{i1}x_{i2}x_{i3}\text{-space}.$$

(ii) Assume that f is of type III with $s_C(-a) = 0$ (4.7(ii)). Moreover we assume that $Q \in C$ has the local description in (4.9(i)). Then $a_1 = w_{Q_1}^C(1) = 1$ and $(H \cdot C) = 1/r$. In particular we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, a)$ of the index one cover $Q^\sharp \in Y^\sharp$ so that

$$Q^\sharp \in C^\sharp \subset Y^\sharp \cong o \in (x_1\text{-axis}) \subset x_1x_2x_3\text{-space}.$$

Proof. (i) First of all by (4.10(i)), we can choose $(A, B) = (1, 2)$ or $(2, 1)$ so that the description at Q_A is of form $(t^{c_A/r_A}, 0, t^{\overline{ac_A}/r_A})$ and that at Q_B is of form $(0, t^{c_B/r_B}, t^{1-\overline{ac_B}/r_B})$.

From the map $\mathcal{O}_Y(-E)^{\otimes r_A r_B} \otimes \mathcal{O}_Y(r_A r_B E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we obtain $r_A r_B s_C(-1) + r_A r_B (E \cdot C) = -r_B w_{Q_A}^C(1) - r_A w_{Q_B}^C(1)$. By $s_C(-1) = -1$ (4.2(iii)), $w_{Q_A}^C(1) = c_A$ and $w_{Q_B}^C(1) = r_B - c_B$, we have

$$(4.11.1) \quad 1 + (H \cdot C) = \frac{c_A}{r_A} + \frac{r_B - c_B}{r_B}.$$

On the other hand from the map $\mathcal{O}_Y(-aE)^{\otimes r_A r_B} \otimes \mathcal{O}_Y(ar_A r_B E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we obtain $r_A r_B s_C(-a) + ar_A r_B (E \cdot C) = -r_B w_{Q_A}^C(a) - r_A w_{Q_B}^C(a)$. By $s_C(-a) = -1$, $w_{Q_A}^C(a) = \overline{ac_A}$ and $w_{Q_B}^C(a) = r_B - \overline{ac_B}$, we have

$$(4.11.2) \quad 1 + a(H \cdot C) = \frac{\overline{ac_A}}{r_A} + \frac{r_B - \overline{ac_B}}{r_B}.$$

From these equalities (4.11.1) and (4.11.2), one of the inequalities $c_A < \overline{ac_A}$ and $c_B > \overline{ac_B}$ holds. However $c_B > \overline{ac_B}$ cannot occur because $(r_B, 0)$ is contained in the semigroup (3.12.1) for Q_B . Hence we have $c_A < \overline{ac_A}$.

We claim that $c_A = 1$. Suppose that $c_A \geq 2$ and take the smallest integer m such that $c_A m \geq r_A$. $w_{Q_A}^C(m) \equiv c_A m - r_A$ modulo r_A and $c_A m - r_A < a_{A1} < a_{A3}$. Hence the map $\mathcal{O}_Y(-mE) \otimes \mathcal{O}_Y(mE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ at Q_A is properly contained in $\mathfrak{m}_{Q_A \subset C}$. In our situation the image of the map $\mathcal{O}_Y(aE) \otimes \mathcal{O}_Y(-aE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ is $\mathfrak{m}_{\{Q_1, Q_2\} \subset C}$. Considering (4.2(iii)) also, we have

(i) $m > a$ or,

(ii) $r_1 \mid m$ and $m < a$.

When (i) occurs, $\overline{ac_A} = ac_A$. Then $c_A = 1$ since $(r_A, 0)$ is contained in the semigroup (3.12.1) for Q_A . When (ii) occurs, unless $m = 2$ the map $\mathcal{O}_Y(-(m-1)E) \otimes \mathcal{O}_Y((m-1)E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ at Q_A is also properly contained in $\mathfrak{m}_{Q_A \subset C}$, which is a contradiction. Thus the case where $m = 2$ in (ii) remains. In this case $r_1 = 2$ and $c_1 = 1$. Then we have $s_C(-a) = 0$ as in the proof of (4.10), which contradicts the assumption $s_C(-a) = -1$.

Now we have $(A, B) = (1, 2)$ and $r_1 < r_2$ by substituting $c_A = 1$ in (4.11.1). We have

$$(4.11.3) \quad (H \cdot C) = \frac{1}{r_1} - \frac{c_2}{r_2},$$

$$(4.11.4) \quad a(H \cdot C) = \frac{\overline{a}}{r_1} - \frac{\overline{ac_2}}{r_2}.$$

Especially $\lfloor a/r_1 \rfloor = \lfloor ac_2/r_2 \rfloor$.

We claim that $c_2 = 1$ and hence $a < r_1$ (3.1.2) and $(H \cdot C) = 1/r_1 - 1/r_2$ (4.11.3). We can show this similarly as we showed $c_1 = 1$ once we obtain $c_2 < r_2 - \overline{ac_2}$. But this inequality comes from the following obtained by (4.11.3) and (4.11.4) respectively.

$$1 - (H \cdot C) = \frac{r_1 - 1}{r_1} + \frac{c_2}{r_2},$$

$$1 + a(H \cdot C) = \frac{\overline{a}}{r_1} + \frac{r_2 - \overline{ac_2}}{r_2}.$$

(ii) By (4.10(ii)) the description at Q is of form $(t^{c/r}, 0, t^{\overline{ac}/r})$ (4.9). From the map $\mathcal{O}_Y(-E)^{\otimes r} \otimes \mathcal{O}_Y(rE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we obtain $rs_C(-1) + r(E \cdot C) = -w_Q^C(1)$. By $s_C(-1) = 0$ (4.2(iv)) and $w_Q^C(1) = c$, we have $(H \cdot C) = c/r$. On the other hand from the map $\mathcal{O}_Y(-aE)^{\otimes r} \otimes \mathcal{O}_Y(arE) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$, we obtain $rs_C(-a) + ar(E \cdot C) = -w_Q^C(a)$. By $s_C(-a) = 0$ and $w_Q^C(a) = \overline{ac}$, we have $a(H \cdot C) = \overline{ac}/r$. Hence we have $ac = \overline{ac}$, and we have the result because $(r, 0)$ is contained in the semigroup (3.12.1). $\square$

The next lemma is an easy consequence by numerical argument.

Lemma 4.12. *Assume that f is of type $\text{IIb}^{\vee\vee}$. Then $(H \cap E)_{\text{red}}$ is irreducible.*

Proof. We may assume that $s_C(-a) = -1$ by (4.8). Write $[H \cap E] = u[C] + [F]$ cycle theoretically, where the support of $[F]$ does not pass Q_1 (4.1(iii)). Then we can write $(H \cdot [H \cap E]) = u(H \cdot C) + v/r_2$. By $(H \cdot [H \cap E]) = E^3 = a^{-1}(1/r_1 + 1/r_2)$ (3.2) and $(H \cdot C) = 1/r_1 - 1/r_2$ (4.11(i)), we obtain

$$\frac{au - 1}{r_1} = \frac{au + 1 - av}{r_2}.$$

Since $r_1 < r_2$ (4.11(i)) and $a \geq 2$, we have $v = 0$. $\square$

4.13. We can take suitable surfaces by (4.4(i)).

- (i) Assume that f is of type $\text{IIb}^{\vee\vee}$. Then by (3.5.1(ii)) and (4.4(i)) there exist surfaces S, S_0 on Y defined by elements in $\mathcal{O}_Y(-aE)$ such that $(S \cap E)_{\text{red}} \not\subseteq (H \cap E)_{\text{red}} \cup (S_0 \cap E)_{\text{red}}$. Let D be an irreducible reduced curve in $S \cap E$ which intersects H and S_0 properly. We can see that $D \cong \mathbb{P}^1$ as in (3.10).
- (ii) Assume that f is of type III. Then by (3.5.1(ii)) and (4.4(i)) we can take a suitable H and a suitable $C \cong \mathbb{P}^1 \subseteq H \cap E$ so that C properly intersects the strict transform H_0 of a general hyperplane section on $P \in X$.

Lemma 4.14. *Assume that f is of type $\text{IIb}^{\vee\vee}$ with $s_C(-a) = -1$, and take $C \subseteq H \cap E$ and $D \subseteq S \cap E$ as in (4.7(i)) and (4.13(i)). Then we can write $[S \cap E] = x[C] + [D]$ cycle theoretically with a positive integer x , and D intersects C exactly at one of Q_1 and Q_2 . Let Q_i ($i = 1$ or 2) be the one D passes through. Then $(H \cdot D) = 1/r_i$ and we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, a)$ of the index one cover $Q_i^\# \in Y^\#$ so that*

$$Q_i^\# \in D^\# \subset Y^\# \cong o \in (x_1\text{-axis}) \subset x_1x_2x_3\text{-space}.$$

Proof. Write $[S \cap E] = x[C] + [D']$ cycle theoretically. We can write $1/r_1 + 1/r_2 = (H \cdot [S \cap E]) = x(H \cdot C) + y/r_1 + z/r_2$. If $y = z = 1$ then $[S \cap E] = [D']$ and $(S \cdot C) \geq 1/r_1 + 1/r_2$ because S passes through Q_1 and Q_2 . On the other hand $(S \cdot C) \leq (S \cdot [H \cap E]) = 1/r_1 + 1/r_2$ (3.2). Hence $C = H \cap E$ and $s_C(-a) \geq 0$ by (3.5.1(ii)) and (4.4(ii)) with $i = a < r_1$ (4.11(i)). It contradicts $s_C(-a) = -1$.

Hence $y = 0$ or $(y, z) = (1, 0)$. By $(H \cdot [S \cap E]) = 1/r_1 + 1/r_2$ and $(H \cdot C) = 1/r_1 - 1/r_2$ (4.11(i)), we obtain

$$\frac{x - 1 + y}{r_1} = \frac{x + 1 - z}{r_2}.$$

Since $y = 0$ or $(y, z) = (1, 0)$, and $r_1 < r_2$ (4.11(i)), we have $(y, z) = (1, 0)$ or $(0, 1)$. Therefore, provided that $r_1 \nmid r_2$, then $D' = D$ and D intersects C only at Q_1 or Q_2 with $(H \cdot D) = 1/r_1, 1/r_2$ respectively. But if $r_1 \mid r_2$ and $(H \cdot D') = 1/r_1$, then setting $(r_1, r_2) = (r, sr)$, by (4.12) we can write

$$au\left(\frac{1}{r} - \frac{1}{sr}\right) = \frac{1}{r} + \frac{1}{sr},$$

where $[H \cap E] = u[C]$ ($u \geq 2$). By this equation and $s > 1$ (4.11(i)), we have $(s, au) = (2, 3)$ or $(3, 2)$. But this contradicts $a \geq 2$ and $u \geq 2$.

Let Q_i ($i = 1$ or 2) be the one D passes through. From the map $\mathcal{O}_Y(-E)^{\otimes r_i} \otimes \mathcal{O}_Y(r_i E) \otimes \mathcal{O}_D \rightarrow \mathcal{O}_D$, we obtain $r_i s_D(-1) + r_i(E \cdot D) = -w_{Q_i}^D(1)$. Hence $w_{Q_i}^D(1) \equiv 1$ modulo r_i . Therefore $w_{Q_i}^D(1) = 1$ since $(r_i, 0)$ is contained in the semigroup (3.12.1) for $Q_i \in D$, whence we have the desired description. $\square$

The next theorem gives complete description in types $\text{Ib}^{\vee\vee}$ and III.

Theorem 4.15. *Assume that f is of type $\text{Ib}^{\vee\vee}$ and III. Then there exist an H and a curve $C \subseteq H \cap E$ in (4.7) which satisfy the assumption in (4.8).*

Proof. We first treat type $\text{Ib}^{\vee\vee}$. Suppose that C does not satisfy the assumption in (4.8), that is, suppose that $s_C(-a) = -1$ (4.2(iii)). Then by (4.13(i)) and (4.14), there exist an S and a $Q_i \in D \cong \mathbb{P}^1$ in (4.13(i)) such that we may choose semi-invariant local coordinates x_1, x_2, x_3 so that $(x_1, x_2, x_3)|_{D^\dagger} = (t^{1/r_i}, 0, 0)$. D intersects H properly.

Let $H^\sharp$ be the preimage of H on the index one cover $Q_i^\sharp \in Y^\sharp$. Write the defining equation of $H^\sharp$ at $Q_i^\sharp$ as $h(x_1) + g_1 x_2 + g_2 x_3$, where $h(x_1)$ is a semi-invariant formal series of weight 1 and g_1, g_2 are semi-invariant formal series of x_1, x_2, x_3 . The order of the equation of $H^\sharp|_{D^\dagger}$ with respect to t equals $(H \cdot D) = 1/r_i$ (4.14). It means that the order of $h(t^{1/r_i})$ with respect to t is $1/r_i$, whence the coefficient of x_1 in h is not zero. Hence we may choose the coordinates x_1, x_2, x_3 so that $H^\sharp$ is given by $x_1 = 0$. If $Q_i = Q_1$ then for the coordinates x_{11}, x_{12}, x_{13} in (4.11(i)) we can write $x_1 = cx_{11} + g(x_{11}, x_{12}, x_{13})$, where c is not zero and g is a semi-invariant formal series of weight 1 which does not contain the term x_{11} . Thus we can easily see that $x_1|_{C^\dagger} \neq 0$ by $(x_{11}, x_{12}, x_{13})|_{C^\dagger} = (t^{1/r_1}, 0, 0)$, which contradicts $C \subset H$.

Hence $Q_i = Q_2$. We consider a general S_0 in (4.13(i)). $S_0 \cap E$ also has a component $D_0 \cong \mathbb{P}^1$ which has the same properties as $D \subset S \cap E$ has, and we can choose new coordinates x_1', x_2', x_3' and t' for $Q_2 \in D_0 \subset Y$ so

that $(x_1', x_2', x_3')|_{D_0^\dagger} = (t'^{1/r_2}, 0, 0)$. Since $(S \cdot D_0) = a(H \cdot D_0) = a/r_2 < 1$ (4.14)(3.1.2), S intersects D_0 only at Q_2 . Let $S^\sharp$ be the preimage of S on $Y^\sharp$, and let $g(x_1, x_2, x_3) = h(x_1', x_2', x_3')$ be the defining equation of $S^\sharp$ at $Q_2^\sharp$. The order of $h(t'^{1/r_2}, 0, 0)$, the equation of $S^\sharp|_{D_0^\dagger}$, with respect to t equals $(S \cdot D_0) = a/r_2$, whence we can write $h = (c' + h_1(x_1')x_1'^{r_2})x_1'^a + h_2(x_1', x_2', x_3')x_2' + h_3(x_1', x_2', x_3')x_3'$, where $c' \in \mathbb{C}^\times$ and h_1, h_2, h_3 are semi-invariant with weights $0, a+1, 0$. Thus we can write $g = cx_1^a + \dots$ ($c \in \mathbb{C}^\times$) and we see that $g(t^{1/r_2}, 0, 0) = 0$ holds only if h_3 is a unit, by considering weights and $a < r_1 < r_2$ (4.13(i)). Of course $g(t^{1/r_2}, 0, 0) = 0$ by $D \subset S$, whence h_3 is a unit and $S^\sharp$ is smooth at $Q_2^\sharp$.

Therefore we may choose the coordinates x_{21}, x_{22}, x_{23} in (4.11(i)) so that $x_{21} = x_1$, the equation of $H^\sharp$, and that x_{23} is the equation of $S^\sharp$. Especially the scheme $S \cap H$ is irreducible and reduced at the generic point of C . Write $[H \cap E] = u[C]$ (4.12) and $[S \cap E] = x[C] + [D]$ (4.14). Then we have

$$\begin{aligned} ua\left(\frac{1}{r_1} - \frac{1}{r_2}\right) &= \frac{1}{r_1} + \frac{1}{r_2}, \\ x\left(\frac{1}{r_1} - \frac{1}{r_2}\right) + \frac{1}{r_2} &= \frac{1}{r_1} + \frac{1}{r_2}. \end{aligned}$$

Thus $(ua - x)(r_2 - r_1) = r_1$ and we can write $r_1 = nr$, $r_2 = (n+1)r$. We note that

$$(4.15.1) \quad ua = 2n + 1,$$

$$(4.15.2) \quad x = n + 1.$$

On the other hand let $\pi: \hat{E} \rightarrow E$ be the normalisation of E and set $\hat{C} = (C \times_E \hat{E})_{\text{red}}$. The coefficients of $[C]$ in the 1-cycles $[S \cap E]$ and $[aH \cap E]$ are determined by the lengths of the schemes $\pi^*S \cap \hat{C}$ and $\pi^*(aH) \cap \hat{C}$ at the generic points of all the irreducible components of $\hat{C}$. From this point of view, we see that $(S \cap E) \cap (aH \cap E)$ contains 1-cycle $x[C]$ because of the general choice of S . $(S \cap E) \cap (aH \cap E) \subseteq S \cap aH$ and $S \cap H$ is irreducible and reduced at the generic point of C . Hence $x \leq a$ but it contradicts (4.15.1), (4.15.2) and $u \geq 2$.

We now treat type III. We start with $H, Q \in C$ and H_0 in (4.13(ii)). Suppose that C does not satisfy the assumption in (4.8), that is, suppose that $s_C(-a) = 0$. Then by (4.9) and (4.11(ii)) we may choose coordinates x_1, x_2, x_3 so that $(x_1, x_2, x_3)|_{C^\dagger} = (t^{1/r}, 0, 0)$, $(t^{2/r}, t^{1-2/r}, 0)$ or $(t^{1-2/r}, t^{2/r}, 0)$. We note that $(H_0 \cdot E) = 1/r, 2/r, 1 - 2/r$ respectively in these cases. Thus H_0 intersects C only at Q .

Let $H_0^\sharp$ be the preimage of H_0 on the index one cover $Q^\sharp \in Y^\sharp$ and let $g(x_1, x_2, x_3)$ be the defining equation of $H_0^\sharp$ at $Q^\sharp$. g is semi-invariant of weight 1 and the order of $g(t^{1/r}, 0, 0)$, $g(t^{2/r}, t^{1-2/r}, 0)$ or $g(t^{1-2/r}, t^{2/r}, 0)$ with respect to t is $1/r, 2/r, 1 - 2/r$ respectively. As in the former argument

we can see that the coefficient of x_1 in the equation of $H_0^\sharp$ is not zero in each of the cases. Hence we may choose semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, a)$ of $Q^\sharp \in Y^\sharp$ so that H_0 is given by $x_1 = 0$.

Now we take $Q \in C_0 \cong \mathbb{P}^1 \subset H_0 \cap E$. If $s_{C_0}(-a) \neq 1$, then $s_{C_0}(-a) = 0$ and again we may choose new coordinates x_1', x_2', x_3' and t' so that $(x_1', x_2', x_3')|_{C_0^\dagger} = (t'^{1/r}, 0, 0)$, $(t'^{2/r}, t'^{1-2/r}, 0)$ or $(t'^{1-2/r}, t'^{2/r}, 0)$. On the other hand we can write $x_1 = cx_1' + g(x_1', x_2', x_3')$, where c is not zero and g is a semi-invariant formal series of weight 1 which does not contain the term x_1' . Thus we can easily see that $x_1|_{C_0^\dagger} \neq 0$, which contradicts $C_0 \subset H_0$. $\square$

5 Existence of Du Val sections

5.1. In this section we show the existence of Du Val sections in the anticanonical system of Y (2.7) together with providing more information. It is done when f is of type O or I (3.7), and we have nothing to do in type IV (3.2.1). Thus we keep the assumption that f is of type II or III. First we restate results in Section 4.

(i) (type IIa (4.5))

- (a) $a = 2$ and $[H \cap E] = [\mathbb{P}^1] + [\mathbb{P}^1]$. $s_C(-2) = 0$ and $(H \cdot C) = 1/r$. $Q \in C$ is locally expressed by semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, 4)$ of the index one cover $Q^\sharp \in Y^\sharp$ so that $(x_1, x_2, x_3)|_{C^\dagger} = (t^{(r+1)/2r}, 0, t^{2/r})$ ($r \equiv 1$ modulo 4) or $(0, t^{(r-1)/2r}, t^{2/r})$ ($r \equiv 3$ modulo 4).
- (b) $r = 5$, $a = 4$ and $H \cap E = \mathbb{P}^1$. $s_C(-4) = 0$ and $(H \cdot C) = 1/r$. $Q \in C$ is locally expressed by semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, 3)$ of the index one cover $Q^\sharp \in Y^\sharp$ as $(x_1, x_2, x_3)|_{C^\dagger} = (t^{3/5}, t^{2/5}, 0)$.

(ii) (type IIb $^\vee$ (4.6))

- (a) $J = \{(r, 1), (r, 1)\}$, $a = 2$ and $H \cap E = \mathbb{P}^1$. $s_C(-2) = 0$ and $(H \cdot C) = 1/r$. $Q \in C$ is locally expressed by semi-invariant local coordinates x_1, x_2, x_3, x_4 with weights $\text{wt}(x_1, x_2, x_3, x_4) = (1, -1, 2, 0)$ of the index one cover $Q^\sharp \in Y^\sharp$ as $(x_1, x_2, x_3, x_4)|_{C^\dagger} = (0, 0, t^{2/r}, t)$.
- (b) $J = \{(2, 1), (4, 1)\}$, $a = 3$ and $H \cap E = \mathbb{P}^1$. $s_C(-3) = 0$ and $(H \cdot C) = 1/4$. $Q \in C$ is locally expressed by semi-invariant local coordinates x_1, x_2, x_3, x_4 with weights $\text{wt}(x_1, x_2, x_3, x_4) = (1, 3, 3, 2)$ of the index one cover $Q^\sharp \in Y^\sharp$ as $(x_1, x_2, x_3, x_4)|_{C^\dagger} = (0, 0, t^{3/4}, t^{1/2})$.

- (iii) (type IIb^{vv} (4.15)) $H \cap E = \mathbb{P}^1$. $s_C(-a) = 0$ and $(H \cdot C) = (r_1 + r_2)/ar_1r_2$. $Q_i \in C$ is locally expressed by semi-invariant local coordinates x_{i1}, x_{i2}, x_{i3} with weights $\text{wt}(x_{i1}, x_{i2}, x_{i3}) = (1, -1, a)$ of the index one cover $Q_i^\# \in Y^\#$ as $(x_{i1}, x_{i2}, x_{i3})|_{C^\dagger} = (0, 0, t^{1/r_i})$.
- (iv) (type III (4.15)) We can choose H so that $H \cap E = \mathbb{P}^1$. $s_C(-a) = 1$ and $(H \cdot C) = (r+1)/ar$. $Q \in C$ is locally expressed by semi-invariant local coordinates x_1, x_2, x_3 with weights $\text{wt}(x_1, x_2, x_3) = (1, -1, a)$ of the index one cover $Q^\# \in Y^\#$ as $(x_1, x_2, x_3)|_{C^\dagger} = (0, 0, t^{1/r})$.

5.2. Let S be a surface on Y defined by a general element in $\mathcal{O}_Y(-aE)$ and let S_X be its strict transform on X . Our main goal is to see that S has at worst Du Val singularities, but we furthermore induce some information on $S \cap E$.

Lemma 5.3. *Assume that f is of type IIa with $a = 2$, and write $[H \cap E] = [C] + [C']$ cycle theoretically. Then there exists a natural exact sequence*

$$0 \rightarrow \mathcal{O}_{C'}(-1) \rightarrow \mathcal{O}_{H \cap E} \rightarrow \mathcal{O}_C \rightarrow 0.$$

Proof. Let $\mathcal{I}$ be the kernel of the natural map $\mathcal{O}_{H \cap E} \rightarrow \mathcal{O}_C$. $\mathcal{I}$ is supported on $C' \cong \mathbb{P}^1$. By $H^0(\mathcal{O}_{H \cap E}) = \mathbb{C}$ (3.10.1), we have $H^0(\mathcal{I}) = 0$. By $H^1(\mathcal{O}_{H \cap E}) = 0$ (3.10(ii)) we have $H^1(\mathcal{I}) = 0$, whence $\mathcal{I} \cong \mathcal{O}_{\mathbb{P}^1}(-1)$. $\square$

Theorem 5.4. (i) *If f is of type II, then S intersects $H \cap E$ at and only at the non-Gorenstein points of Y .*

(ii) *If f is of type III, then for any $Q' \in C = H \cap E$ which is a smooth point of Y , there exists an S such that S intersects $H \cap E$ at and only at Q' and the non-Gorenstein point of Y .*

Proof. (i) Take any $C \cong \mathbb{P}^1 \subseteq H \cap E$. By (3.4.3), (3.9) and (5.3) we obtain a surjective map $f_*\mathcal{O}_Y(-aE) \rightarrow H^0([\mathcal{O}_Y(-aE)]_C)$. Recalling that $s_C(-a) = 0$ (5.1), we have $H^0([\mathcal{O}_Y(-aE)]_C) = \mathbb{C}$. Hence we find an S which properly intersects $H \cap E$. Of course S passes through any non-Gorenstein point of Y . On the other hand because $(S \cdot [H \cap E]) = aE^3 \leq 1$ in any case (5.1), we see that S does not intersect $H \cap E$ at any other point of Y .

(ii) It suffices to consider $[\mathcal{O}_Y(-aE) \otimes \mathfrak{m}_{Q' \subset Y}]_C$ instead of $[\mathcal{O}_Y(-aE)]_C$ since $s_C(-a) = 1$ (5.1). $\square$

It is the time to show our main theorem (2.7).

Theorem 5.5. *Assume that f is of type II or III. Then for a suitable S in (5.4), S has at worst Du Val singularities. Moreover the type of a Du Val singularity at any $Q \in S$ is that of a general Du Val section of a germ $Q \in Y$.*

Proof. We have only to consider the case where P is a singular point by [Ka1]. Because of (5.4) and Bertini's theorem, it is enough to show that S has a Du Val singularity at each non-Gorenstein point $Q \in Y$ whose type equals that of a general Du Val section of a germ $Q \in Y$.

First we treat type IIa (5.1(i)). Let $Q \in C = H \cap E$ be the non-Gorenstein point with a curve investigated.

Consider the case $a = 2$. It is enough to show that the coefficient of x_3 in the semi-invariant equation $h(x_1, x_2, x_3)$ of S in (5.2) is not zero. Since S intersects C only at Q (5.4), the order of $h(t^{(r+1)/2r}, 0, t^{2/r})$ ($r \equiv 1$ modulo 4) or $h(0, t^{(r-1)/2r}, t^{2/r})$ ($r \equiv 3$ modulo 4) with respect to t equals $(S \cdot C) = 2/r$. Hence the coefficient of x_3 in h is not zero.

Consider the case $a = 4$. $-K_Y$ is linearly equivalent to $E + 5H$. Thus it is enough to show that

- (i) E has a Du Val singularity of type A_4 at Q , and
- (ii) $|5H|$ is free at Q .

To see (i), consider a birational morphism $g: Z \rightarrow Y$ such that Z has a g -exceptional divisor F whose discrepancy with respect to K_X is 1 [Ma]. Write

$$\begin{aligned} K_Z &= g^*K_Y + bF + (\text{others}), \\ g^*E &= E_Z + mF + (\text{others}), \end{aligned}$$

where E_Z is the strict transform of E . Then $K_Z = g^*(f^*K_X + 4E) + bF + (\text{others}) = g^*f^*K_X + 4E_Z + (b + 4m)F + (\text{others})$, whence $b = m = 1/5$ by $5b, 5m \in \mathbb{Z}$. Hence the defining equation of E by x_1, x_2, x_3 in (5.1(i)) has a nonzero linear term. Considering the weight we see that the coefficient of x_3 in the equation of E is not zero, whence we may take x_3 so that E is given by $x_3 = 0$. This implies (i).

It is easy to see (ii). From (3.9) and (3.4.3) we obtain a surjective map $f_*\mathcal{O}_Y(-5E) \twoheadrightarrow H^0([\mathcal{O}_Y(-5E)]_C) = H^0(\mathcal{O}_{\mathbb{P}^1}(1))$, which shows (ii).

Secondly we treat type IIb^v (5.1(ii)). Let $Q \in C = H \cap E$ be the non-Gorenstein point with a curve investigated. $Q \in Y$ is described as in (3.2.2). Consider a surface T on a germ $Q \in Y$ defined by a section in $\mathcal{O}_Y(-K_Y)_Q$ whose equation in y_1, y_2, y_3, y_4 contains a linear term. Then by the list (3.2.2) we have

- (i) If $J = \{(r, 1), (r, 1)\}$ and $r \geq 5$, then the linear term of the equation of T is y_3 . Any such T gives a Du Val singularity of type A_{2r-1} , which is the type of a general Du Val section of a germ $Q \in Y$.
- (ii) In any other case, the linear term of the equation of T is a linear combination of y_2 or y_3 . Any T equipped with a general linear term gives a Du Val singularity of type A_5, E_6 or D_5 according to (ia), (ib)

or (ii) in (3.2.2) respectively, which is the type of a general Du Val section of a germ $Q \in Y$.

Thus if $J = \{(r, 1), (r, 1)\}$ and $r \geq 5$, it is enough to show that

5.5.1. The equation of S in (5.2) by y_1, y_2, y_3, y_4 contains a linear term.

In any other case, $-K_Y$ is linearly equivalent to $E + rH$. Thus it is enough to show that, adding to (5.5.1),

5.5.2. (i) The equation of E by y_1, y_2, y_3, y_4 contains a linear term which is linearly independent to that in the equation of S .

(ii) $|rH|$ is free at Q .

However in (5.5.2), the existence of a linear term in the equation of E and the statement (5.5.2(ii)) can be shown similarly as in the proof of type IIa with $a = 4$. In particular in (5.5.2) we may choose coordinates x_1, x_2, x_3, x_4 in (5.1(ii)) so that E is given by $x_2 = 0$. Hence (5.5.1) and (5.5.2) follows if we prove that the coefficient of x_3 in the defining equation $h(x_1, x_2, x_3, x_4)$ of S in (5.2) is not zero.

Because S intersects C only at Q (5.4), the order of $h(0, 0, t^{2/r}, t)$ or $h(0, 0, t^{3/4}, t^{1/2})$ with respect to t equals $(S \cdot C) = aE^3 = 2/r$ or $3/4$ respectively according to $J = \{(r, 1), (r, 1)\}$ or $\{(2, 1), (4, 1)\}$. Hence the coefficient of x_3 in h is not zero.

Finally we treat types IIb^{vv} and III (5.1(iii)(iv)). Let $Q \in C = H \cap E$ be a non-Gorenstein point with a curve investigated. It is enough to show that the coefficient of x_3 in the defining equation $h(x_1, x_2, x_3)$ of S in (5.2) is not zero, and we can see this similarly, remarking that the local intersection number of S and C at Q is $1/r_Q$ (5.1(iii)(iv))(5.4). $\square$

6 Possible types of singularities

6.1. In this section we restrict types of Du Val singularities on S and S_X , and prove (2.8) as a consequence. First we recall dual graphs for minimal resolutions of Du Val singularities. $\circ$ denotes an exceptional curve, $\bullet$ denotes the strict transform of a general hyperplane section, and numbers attached exceptional curves F_i are the coefficients of F_i in the fundamental cycle.

(i) (type A_n)

$$\bullet - \frac{F_1}{\circ} - \dots - \frac{F_n}{\circ} - \bullet$$

(ii) (type D_n)

$$\begin{array}{ccccccc}
 & & \bullet & & & & F_n \\
 & & | & & & & \circ \\
 & & 1 & & & & 1 \\
 & & | & & & & | \\
 F_1 & - & F_2 & - & \dots & - & F_{n-2} & - & F_{n-1} \\
 \circ & & \circ & & & & \circ & & \circ \\
 1 & & 2 & & & & 2 & & 1
 \end{array}$$

(iii) (type E_6)

$$\begin{array}{ccccccc}
 & & & & \bullet & & \\
 & & & & | & & \\
 & & & & F_1 & & \\
 & & & & \circ & & \\
 & & & & 2 & & \\
 & & & & | & & \\
 F_2 & - & F_3 & - & F_4 & - & F_5 & - & F_6 \\
 \circ & & \circ & & \circ & & \circ & & \circ \\
 1 & & 2 & & 3 & & 2 & & 1
 \end{array}$$

(iv) (type E_7)

$$\begin{array}{ccccccc}
 & & & & F_7 & & \\
 & & & & \circ & & \\
 & & & & 2 & & \\
 & & & & | & & \\
 \bullet & - & F_1 & - & F_2 & - & F_3 & - & F_4 & - & F_5 & - & F_6 \\
 & & \circ & & \circ & & \circ & & \circ & & \circ & & \circ \\
 & & 2 & & 3 & & 4 & & 3 & & 2 & & 1
 \end{array}$$

(v) (type E_8)

$$\begin{array}{ccccccc}
 & & & & & & F_8 \\
 & & & & & & \circ \\
 & & & & & & 3 \\
 & & & & & & | \\
 \bullet & - & F_1 & - & F_2 & - & F_3 & - & F_4 & - & F_5 & - & F_6 & - & F_7 \\
 & & \circ & & \circ & & \circ & & \circ & & \circ & & \circ & & \circ \\
 & & 2 & & 3 & & 4 & & 5 & & 6 & & 4 & & 2
 \end{array}$$

6.2. Take a surface S on Y defined by a general element in $|-K_Y|$ (3.8)(5.5), and define S_X as its strict transform on X . A general hyperplane section $P \in H_X$ on X gives also a general hyperplane section $P \in D_X = H_X|_{S_X}$ on S_X . Let $f_S: S \rightarrow S_X$ be the induced map. Since S and S_X have at worst Du Val singularities (3.8)(5.5), f_S factors the minimal resolution of S_X . Let Z be the strict transform on S of the fundamental cycle. $f^*H_X = H + bE$ in (3.7) and $f^*D_X = D + Z = H|_S + bE|_S$, where D is the strict transform on S of D_X . We note that $bE|_S \leq Z$. We also have

$$(6.2.1) \quad (H|_S \cdot bE|_S) = b^2 a E^3.$$

6.3. Assume that f is of type I. In this case one of the following holds by (3.2), (3.5) and (3.8).

- (i) $J = \{(7, 3)\}$, $E^3 = 1/7$ and $a = b = 2$.
- (ii) $J = \{(3, 1), (5, 2)\}$, $E^3 = 1/15$ and $a = b = 2$.

Lemma 6.4. *Assume that f is of type I. Take a general S in (6.2).*

- (i) *If $J = \{(7, 3)\}$ (6.3(i)), then the dual graph for the partial resolution $S \rightarrow S_X$ is*

$$* - \underset{2}{\circ} \quad \text{or} \quad \underset{2}{\circ} - * - \underset{2}{\circ},$$

where the attached numbers are the coefficients in $2E|_S$, and $$ means that a Du Val singularity of type at best A_6 appears there.*

- (ii) *If $J = \{(3, 1), (5, 2)\}$ (6.3(ii)), then the dual graph for the partial resolution $S \rightarrow S_X$ is*

$$* - \underset{2}{\circ} - \star \quad \text{or} \quad * - \underset{4}{\circ} - \star,$$

the attached number is the coefficient in $2E|_S$, and $$, $\star$ means that a Du Val singularity of type at best A_2 , A_4 appears there respectively.*

Proof. (i) Since $(-E \cdot [S \cap E]) = 2E^3 = 2/7$, it is enough to show that the existence of S such that $S \cap E$ defines a reduced 1-cycle. In this case we can calculate $\chi(\mathcal{Q}_{-2}) = 2$ by (3.4.2). Hence we have a desired S as in the proof of (4.4(i)).

(ii) Since $(-E \cdot [S \cap E]) = 2E^3 = 2/15$, it is enough to show that the existence of S such that $S \cap E$ defines an irreducible but possibly nonreduced 1-cycle. By $(-E \cdot [S \cap E]) < 1/5$, each irreducible component of $(S \cap E)_{\text{red}}$ passes through all the non-Gorenstein points of Y . Hence $(S \cap E)_{\text{red}} = \mathbb{P}^1$ by $h^1(\mathcal{O}_{S \cap E}) = 0$ (3.10.1). $\square$

6.5. We restrict the dual graph for the partial resolution S in (6.2) considering $2E|_S \leq Z$ and (6.4). The following list shows all the possibilities up to permutation in terms of F_i in (6.1).

$J = \{(7, 3)\}$ (6.3(i))			$J = \{(3, 1), (5, 2)\}$ (6.3(ii))		
case	S_X	Z	case	S_X	Z
(a)	D_n	$Z \not\supseteq F_1, \dots, F_{m-1}$ and $Z \supseteq 2F_m$	(a)	D_n	$Z \not\supseteq F_1, \dots, F_{m-1}$ and $Z \supseteq 2F_m$
(b)	E_7	$Z \supseteq 2F_1$	(b)	E_7	$Z = 3F_4$
(c)	E_7	$Z = 2F_5$	(c)	E_8	$Z = 4F_3$
(d)	E_7	$Z = 2F_7$	(d)	E_8	$Z = 5F_4$
(e)	E_8	$Z \supseteq 2F_1$	(e)	E_8	$Z = 6F_5$
(f)	E_8	$Z = 3F_2$			
(g)	E_8	$Z = 3F_2 + 2F_7$			
(h)	E_8	$Z = 4F_3$			
(i)	E_8	$Z = 4F_3 + 2F_7$			
(j)	E_8	$Z = 4F_6$			
(k)	E_8	$Z = 2F_7$			
(l)	E_8	$Z = 2F_7 + 3F_8$			
(m)	E_8	$Z = 3F_8$			

We calculate $(H|_S \cdot 2E|_S)$ for all the possible cases.

$J = \{(7, 3)\}$			$J = \{(3, 1), (5, 2)\}$		
case	$2E_S$	$(H _S \cdot 2E _S)$	case	$2E_S$	$(H _S \cdot 2E _S)$
(a)	$2F_m + \dots$	$4/m$	(a)	$2F_m$	$4/m$
(b)	$2F_1 + \dots$	2	(b)	$2F_4$	$8/15$
(c)	$2F_5$	1	(c)	$2F_3$	$1/3$
(d)	$2F_7$	$8/7$	(c)	$4F_3$	$4/3$
(e)	$2F_1 + \dots$	2	(d)	$2F_4$	$1/5$
(f)	$2F_2$	$2/3$	(d)	$4F_4$	$4/5$
(g)	$2F_2 + 2F_7$	1	(e)	$2F_5$	$2/15$
(h)	$2F_3$	$1/3$	(e)	$4F_5$	$8/15$
(i)	$2F_3 + 2F_7$	$4/3$			
(j)	$2F_6$	$2/7$			
(k)	$2F_7$	1			
(l)	$2F_7 + 2F_8$	$8/7$			
(m)	$2F_8$	$1/2$			

$(H|_S \cdot 2E|_S) = 8E^3$ (6.2.1), which is $8/7$ (6.3(i)), $8/15$ (6.3(ii)) respectively. Thus we have the following.

Theorem 6.6. *If f is of type I then P is cE_7 or cE_8 .*

Remark 6.6.1. Actually P is cE_7 when $J = \{(7, 3)\}$. By (6.5) it suffices to exclude the case where $P \in S_X$ is of type E_8 and $Z = 2F_7 + 3F_8$. In the case $S \cap E$ defines the 1-cycle $F_7 + F_8$ on S , and $(E \cdot F_7), (E \cdot F_8)$ must be $-1/7$. On the other hand using the dual graph for the minimal resolution (6.1) we obtain $(E|_S \cdot F_7) = (F_7 + F_8 \cdot F_7)_S = -3/7$ and $(E|_S \cdot F_8) = (F_7 + F_8 \cdot F_8)_S = 1/7$. It is a contradiction.

Example 6.7. There exist examples of type I. The weighted blowup of $o \in (x_1^2 + x_2^3 + x_2x_3^3 + x_4^7 = 0) \subset \mathbb{C}^4$ with weights $\text{wt}(x_1, x_2, x_3, x_4) = (7, 5, 3, 2)$ is an example of type I whose P is cE_7 , and the weighted blowup of $o \in (x_1^2 + x_2^3 + x_3^5 + x_4^7 = 0) \subset \mathbb{C}^4$ with weights $\text{wt}(x_1, x_2, x_3, x_4) = (7, 5, 3, 2)$ is an example of type I whose P is cE_8 .

6.8. On the other hand if f is of type II or III, then by (5.4), (5.5) and (3.2.2), we obtain the dual graph for the partial resolution $S \rightarrow S_X$ below.

(i) (type IIa or $\text{IIb}^\vee$) The dual graph is

$$\bullet - * \begin{cases} \circ \\ 1 \\ \vdots \\ \circ \\ 1 \end{cases}$$

where $\bullet$ denotes $H|_S = D$, the attached numbers are the coefficients in $E|_S = Z$, and $*$ means that a Du Val singularity of type

- (a) A_{r-1} if f is of type IIa,
- (b) A_{2r-1} if f is of type $\text{IIb}^\vee$ with $J = \{(r, 1), (r, 1)\}$, $r \geq 5$,
- (c) A_5 or E_6 if f is of type $\text{IIb}^\vee$ with $J = \{(3, 1), (3, 1)\}$,
- (d) D_5 if f is of type $\text{IIb}^\vee$ with $J = \{(2, 1), (4, 1)\}$,

appears there respectively. The number of exceptional curves $\circ$ is at most 4 in type IIa and at most a in type $\text{IIb}^\vee$. This bound comes from the value of $(H \cdot [S \cap E]) = aE^3$.

(ii) (type $\text{IIb}^{\vee\vee}$ or III) Set $r_1 = 1$, $r_2 = r$ if f is of type III. $P \in S_X$ is of type A_n because $H|_S$ intersects $E|_S$ at two points. The dual graph is

$$\bullet - * - \underset{1}{\circ} - \star - \bullet \quad \text{or} \quad \bullet - * - \underset{1}{\circ} - \underset{1}{\circ} - \star - \bullet,$$

where $\bullet$ denotes $H|_S = D$, the attached numbers are the coefficients in $E|_S = Z$, and $*$, $\star$ means that a Du Val singularity of type A_{r_1-1} , A_{r_2-1} appears there respectively. Since all the components of $(S \cap E)_{\text{red}}$ are numerically proportional, the number of exceptional curves $\circ$ is one or two.

The following list shows all the possibilities of the partial resolution $S \rightarrow S_X$ up to permutation in terms of F_i in (6.1).

type of f	S_X	Z	$(H _S \cdot E _S)$
IIa	D_r	F_r	$4/r$
IIa	D_{r+1}	$F_r + F_{r+1}$	$4/r$
IIa	D_{r+1}	$F_1 + F_{r+1}$	$1 + 1/r$
IIa	D_{r+2}	$F_1 + F_{r+1} + F_{r+2}$	$1 + 1/r$
$\text{IIb}^\vee, J = \{(r, 1), (r, 1)\}$	D_{2r}	F_{2r}	$2/r$
$\text{IIb}^\vee, J = \{(r, 1), (r, 1)\}$	D_{2r+1}	$F_{2r} + F_{2r+1}$	$2/r$
$\text{IIb}^\vee, J = \{(r, 1), (r, 1)\}$	D_{2r+1}	$F_1 + F_{2r+1}$	$1 + 1/2r$
$\text{IIb}^\vee, J = \{(3, 1), (3, 1)\}$	E_7	F_6	$2/3$
$\text{IIb}^\vee, J = \{(2, 1), (4, 1)\}$	D_6	F_1	1
$\text{IIb}^\vee, J = \{(2, 1), (4, 1)\}$	E_6	F_2	$3/4$
$\text{IIb}^{\vee\vee}$	$A_{r_1+r_2-1}$	F_{r_1}	$1/r_1 + 1/r_2$
$\text{IIb}^{\vee\vee}$	$A_{r_1+r_2}$	$F_{r_1} + F_{r_1+1}$	$1/r_1 + 1/r_2$
III	A_r	F_r	$1 + 1/r$
III	A_{r+1}	$F_r + F_{r+1}$	$1 + 1/r$

$(H|_S \cdot E|_S) = aE^3$ (6.2.1). Combining it with (3.2) we have the following theorem.

Theorem 6.9. *Assume that f is of type II or III. Then possible types of Du Val singularities on S and S_X are as follows.*

type of f	type of S_X	type of S
IIa	D_r or D_{r+1}	A_{r-1}
$\text{IIb}^\vee, J = \{(r, 1), (r, 1)\}$	D_{2r} or D_{2r+1}	A_{2r-1}
$\text{IIb}^\vee, J = \{(3, 1), (3, 1)\}$	E_7	E_6
$\text{IIb}^\vee, J = \{(2, 1), (4, 1)\}$	E_6	D_5
$\text{IIb}^{\vee\vee}$	$A_{r_1+r_2-1}$ or $A_{r_1+r_2}$	A_{r_1-1} and A_{r_2-1}
III	A_r or A_{r+1}	A_{r-1}

Furthermore the exceptional locus of the partial resolution $S \rightarrow S_X$ is irreducible unless the type of S_X is that after the word “or” in the above table.

Remark 6.9.1. In types $\text{IIb}^{\vee\vee}$ and III, the type of S_X has to be in fact $A_{r_1+r_2-1}$, A_r respectively by [Ka1], [Ka2] and (2.12) proved in Section 7.

Example 6.10. Let f be the weighted blowup of $o \in (x_1^2 + x_2^2 x_3 + x_3^{2r} + x_4^r = 0) \subset \mathbb{C}^4$ with weights $\text{wt}(x_1, x_2, x_3, x_4) = (r, r, 1, 2)$, where $r \geq 3$ is an odd integer. P is $\text{c}D_{r+1}$ and f is of type $\text{IIb}^\vee$ with $J = \{(r, 1), (r, 1)\}$. The exceptional locus of the partial resolution $S \rightarrow S_X$ is reducible.

Example 6.11. Let f be the weighted blowup of $o \in (x_1^2 + x_2^2 x_3 + x_3^3 + x_4^3 = 0) \subset \mathbb{C}^4$ with weights $\text{wt}(x_1, x_2, x_3, x_4) = (3, 1, 4, 2)$. P is $\text{c}D_4$ and f is of type $\text{IIb}^\vee$ with $J = \{(2, 1), (4, 1)\}$.

7 Divisorial contractions to cA_n points

We begin with introducing a general method to determine f .

Lemma 7.1. *Let $f: (Y \supset E) \rightarrow (X \ni P)$ be a germ of a three dimensional divisorial contraction to a cDV point P and let $K_Y = f^*K_X + aE$. X is identified with a hypersurface in $\bar{X} = \mathbb{C}^4 = x_1x_2x_3x_4$ -space,*

$$P \in X \cong o \in (\phi = 0) \subset \bar{X} = \mathbb{C}^4 = x_1x_2x_3x_4\text{-space}.$$

Let m_i be the multiplicity of $\text{div} x_i$ along E , that is, the largest integer such that $x_i \in f_\mathcal{O}_Y(-m_iE)$. We assume that m_1, m_2, m_3, m_4 have no common factors. Let d be the weighted order of ϕ with respect to weights $\text{wt}(x_1, x_2, x_3, x_4) = (m_1, m_2, m_3, m_4)$ and decompose ϕ as*

$$\phi = \phi_d(x_1, x_2, x_3, x_4) + \phi_{>d}(x_1, x_2, x_3, x_4).$$

where ϕ_d be the weighted homogeneous part of weighted degree d and $\phi_{>d}$ be the part of weighted degree greater than d . Set $c = m_1 + m_2 + m_3 + m_4 - 1 - d$. Let $\bar{g}: (\bar{Z} \supset \bar{F}) \rightarrow (\bar{X} \ni P)$ be the weighted blowup of $\bar{X}$ with weights $\text{wt}(x_1, x_2, x_3, x_4) = (m_1, m_2, m_3, m_4)$, let $\bar{F}$ be its exceptional divisor, and let $\bar{D}_i$ be the strict transform on $\bar{Z}$ of the hyperplane $x_i = 0$ in $\bar{X}$. Let Z be the strict transform of X on $\bar{Z}$, and let $g: Z \rightarrow X$ be the induced morphism. Assume that

- (i) $\bar{F} \cap Z$ defines an irreducible reduced 2-cycle F on Z , and $F \not\subseteq \text{Sing} Z$,
- (ii) $\dim(\text{Sing} \bar{Z} \cap Z) \leq 1$,
- (iii) $\text{Sing} Z \subseteq \bigcup_{1 \leq i \leq 4} \bar{D}_i$, and
- (iv) $F \not\subseteq \bigcup_{1 \leq i \leq 4} \bar{D}_i$.

Then $c \leq a$, and the equality holds if and only if $f \cong g$ over X .

Proof. Z is R_1 from (i), and is Cohen-Macaulay since $Z \subset \bar{Z}$ is locally a cyclic quotient of a hypersurface in $\mathbb{C}^4$. Thus Z is normal. From (ii) we can use the adjunction formula and have $K_Z = (K_{\bar{Z}} + Z)|_Z = g^*K_X + cF$. We note that $-F$ is f -ample and $\mathbb{Q}$ -Cartier.

Consider the centre on Z of the valuation corresponding to E . Because the multiplicity of $\text{div} x_i$ along E equals to that along F , this centre is not contained in $\bigcup_{1 \leq i \leq 4} \bar{D}_i$ by $\bigcap_{1 \leq i \leq 4} \bar{D}_i = \emptyset$ and (iv), whence it intersects the smooth locus of Z by (iii). Thus the discrepancy a of E with respect to K_X is greater than or equal to c , that of F , and the equality holds if and only if $f \cong g$ over X by [Ka1, Lemma 3.4]. $\square$

7.2. Let $f: (Y \supset E) \rightarrow (X \ni P)$ be a germ of a divisorial contraction to a cA_n point P ($n \geq 2$). X is identified with a hypersurface $\phi = 0$ in $\mathbb{C}^4 = x_1x_2x_3x_4$ -space as follows.

$$(7.2.1) \quad P \in X \cong o \in (\phi = x_1x_2 + g(x_3, x_4) = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4\text{-space},$$

where the total order of g with respect to x_3, x_4 is $n + 1$. From (3.2.1) and (6.6), f is of type O, II or III. Our goal in this section is the classification of these contractions (2.12).

7.3. We start with an identification (7.2.1). Set m_i as the multiplicity of $\text{div} x_i$ along E . By coordinates change in which $x_4 \mapsto x_4 + x_i$ ($i = 1, 2$ or 3) if necessary, we may assume that $m_4 = 1$. By coordinates change $x_3 \mapsto x_3 + h(x_4)$ if necessary we may furthermore assume that $x_3 + h(x_4) \notin f_*\mathcal{O}_Y(-(m_3 + 1)E)$ for any h . Decompose ϕ as

$$\phi = \phi_{\leq m_1+m_2}(x_1, x_2, x_3, x_4) + \phi_{>m_1+m_2}(x_1, x_2, x_3, x_4),$$

where $\phi_{\leq m_1+m_2}$ is the part of weighted degree less than or equal to $m_1 + m_2$ and $\phi_{>m_1+m_2}$ is the part of weighted degree greater than $m_1 + m_2$ with respect to weights $\text{wt}(x_1, x_2, x_3, x_4) = (m_1, m_2, m_3, 1)$.

We focus on $\phi_{\leq m_1+m_2} = x_1x_2 + h(x_3, x_4)$. Of course $h \neq 0$. The homogeneous part h_0 in h of the minimal weighted degree d_0 with respect to weights $\text{wt}(x_3, x_4) = (m_3, 1)$ always decomposes into a product of $x_4^{d_0 - m_3 \lfloor \frac{d_0}{m_3} \rfloor}$ and $\lfloor \frac{d_0}{m_3} \rfloor$ linear combinations of x_3 and $x_4^{m_3}$. Hence $h_0 \notin f_*\mathcal{O}_Y(-(d_0 + 1)E)$ because of the assumption $x_3 + h(x_4) \notin f_*\mathcal{O}_Y(-(m_3 + 1)E)$ for any h . Therefore d_0 must be $m_1 + m_2$.

We can see that this choice of coordinates satisfies the assumptions in (7.1), whence

$$a \geq m_1 + m_2 + m_3 + 1 - 1 - (m_1 + m_2) = m_3,$$

and the equality holds if and only if $f \cong g$ in (7.1) over X .

In particular we say the following in type O.

Lemma 7.4. *If f is of type O, then f is the weighted blowup of X with weights $\text{wt}(x_1, x_2, x_3, x_4) = (i, n + 1 - i, 1, 1)$ ($1 \leq i \leq n$) in (7.2.1).*

Lemma 7.5. *f is not of type IIa with $a = 4$.*

Proof. If f is of type IIa with $a = 4$, then $J = \{(5, 2)\}$ from (4.5), whence Y has exactly one non-Gorenstein point and it is a quotient singularity of type $\frac{1}{5}(1, -1, 3)$. Consider a birational morphism $h: Z \rightarrow Y$ such that Z has an h -exceptional divisor F whose discrepancy with respect to K_X is 1. We can

choose n different F as valuations corresponding to exceptional divisors of the weighted blowups of X with weights $\text{wt}(x_1, x_2, x_3, x_4) = (i, n+1-i, 1, 1)$ ($1 \leq i \leq n$) in (7.2.1). Write

$$\begin{aligned} K_Z &= h^*K_Y + bF + (\text{others}), \\ h^*E &= E_Z + mF + (\text{others}), \end{aligned}$$

where E_Z is the strict transform of E . Then $K_Z = h^*(f^*K_X + 4E) + bF + (\text{others}) = h^*f^*K_X + 4E_Z + (b + 4m)F + (\text{others})$, whence $b = m = 1/5$ by $5b, 5m \in \mathbb{Z}$. Hence F is determined uniquely because Y has a quotient singularity of index 5 as the unique non-Gorenstein point. It contradicts that we can choose n different F . $\square$

The case where f is of type IIa with $a = 2$, type IIb or III remains. We can see the next lemma by following [Ka2, Claim 6.13] faithfully.

Lemma 7.6. *Let $f: (Y \supset E) \rightarrow (X \ni P)$ be a germ of a three dimensional divisorial contraction to a singular cDV point P , and let $K_Y = f^*K_X + aE$. Assume that f is of type IIb or III, and set $r_1 = 1$ if f is of type III. $P \in X$ is identified as*

$$P \in X \cong o \in (\phi = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4\text{-space}.$$

Assume that the multiplicity of $\text{div} x_4$ along E is 1. Then

(i) for $1 \leq j \leq \min\{r_1, a\}$,

$$f_*\mathcal{O}_Y(-jE) = (x_{1j}, x_{2j}, x_{3j}, x_4^j)\mathcal{O}_X$$

for some $x_{ij} = x_i + p_{ij}(x_4)$ ($i = 1, 2, 3$), where

$$p_{ij} \in \bigoplus_{k=1}^{j-1} \mathbb{C}x_4^k \subset \mathbb{C}[x_4].$$

(ii) Assume that $r_1 < a$.

(a) The kernel of the linear map

$$\begin{aligned} \mathbb{C}x_{1r_1} \oplus \mathbb{C}x_{2r_1} \oplus \mathbb{C}x_{3r_1} &\rightarrow \\ f_*\mathcal{O}_Y(-r_1E) / (f_*\mathcal{O}_Y(-(r_1+1)E) + (x_4^{r_1})\mathcal{O}_X) & \end{aligned}$$

is of dimension 2.

(b) We assume that $x_{3r_1} \notin f_*\mathcal{O}_Y(-(r_1+1)E) + (x_4^{r_1})\mathcal{O}_X$. Under this situation, for $r_1 < j \leq a$,

$$f_*\mathcal{O}_Y(-jE) = (x_{1j}, x_{2j})\mathcal{O}_X + \sum_{(k,l) \in \bigcup_{s \geq j} J_s} (x_{3r_1}^k x_4^l)\mathcal{O}_X$$

for some $x_{ij} = x_i + p_{ij}(x_{3r_1}, x_4)$ ($i = 1, 2$), where

$$p_{ij} \in \bigoplus_{(k,l) \in \bigcup_{1 \leq s < j} J_s} \mathbb{C}x_{3r_1}^k x_4^l \subset \mathbb{C}[x_{3r_1}, x_4],$$

$$J_s = \{(k, l) \in \mathbb{Z}_{\geq 0}^2 \mid r_1 k + l = s\}.$$

Remark 7.6.1. The similar result holds even if f is of type IIa with $a = 2$ from $d(-1) = \chi(\mathcal{Q}_{-1}) = 1$ (3.5.1(i)). Assume that f is of type IIa with $a = 2$ and the multiplicity of $\text{div} x_4$ along E is 1. Then $x_1, x_2, x_3 \in f_*\mathcal{O}_Y(-2E) + (x_4^2)\mathcal{O}_X$.

7.7. Assume that f is of type IIa with $a = 2$ or type IIb $^\vee$ with $J = \{(r, 1), (r, 1)\}$. Then $m_3 \geq a$ in (7.3) by (7.6), (7.6.1). Assume that f is of type IIb $^{\vee\vee}$ or III, and set $r_1 = 1, r_2 = r$ if f is of type III. We start with an identification (7.2.1). Set m_i as the multiplicity of $\text{div} x_i$ along E . We may assume that $m_4 = 1$ by a coordinate change. Because a general element in $f_*\mathcal{O}_Y(-aE)$ gives a Du Val singularity $P \in S_X$ at worst of type $A_{r_1+r_2}$ (6.9), this element must be of form $cx_3 + \dots$ with $c \in \mathbb{C}^\times$ by (7.6). Hence by coordinates change such that $x_3 \mapsto x_3 + h(x_1, x_4)$ or $x_3 + h(x_2, x_4)$ if necessary we may assume that $x_3 \in f_*\mathcal{O}_Y(-aE)$ (7.6). Therefore $m_3 \geq a$ in (7.3). Hence we have the following from (7.3).

Proposition 7.8. *Assume that f is of type IIb or III. If f is not of type IIb $^\vee$ with $J = \{(2, 1), (4, 1)\}$, then f is the weighted blowup of X with weights $\text{wt}(x_1, x_2, x_3, x_4) = (m_1, m_2, a, 1)$ with respect to the coordinates in (7.3).*

Finally we have to consider the exceptional case where f is of type IIb $^\vee$ with $J = \{(2, 1), (4, 1)\}$.

Lemma 7.9. *If f is of type IIb $^\vee$ with $J = \{(2, 1), (4, 1)\}$, then*

- (i) P is a cA_2 point isomorphic to $o \in (x_1x_2 + x_3^3 + g_{\geq 4}(x_3, x_4) = 0) \subset \mathbb{C}^4 = x_1x_2x_3x_4$ -space, where the total order of $g_{\geq 4}$ with respect to x_3, x_4 is greater than or equal to 4,
- (ii) Y has exactly one non-Gorenstein point Q , which is isomorphic to $o \in (y_1^2 + y_2^2 + y_3^2 + y_4^3 = 0)$ in the quotient space of $\mathbb{C}^4 = y_1y_2y_3y_4$ -space divided by $\mathbb{Z}/(4)$ with weights $\text{wt}(y_1, y_2, y_3, y_4) = (1, 3, 3, 2)$, and
- (iii) $K_Y = f^*K_X + 3E$.

Proof. From (3.2.2), $a = 3$ (iii) and Y has exactly one non-Gorenstein point Q . The germ $Q \in Y$ is described as in (3.2.2(ii)). Because the Gorenstein index r of Y is 4 and $a = 3$, for any valuation v with discrepancy 1 with respect to K_X the centre of v on Y is Q and v has discrepancy $1/4$ with

respect to K_Y , as in the proof of (7.5). Since there are n such valuations as we mentioned in the proof of (7.5), $Q \in Y$ must have an isomorphism in (ii) by the description in (3.2.2(ii)) and [Ha, Theorem 7.4]. In this case there exist exactly two valuations centred at Q with discrepancy $1/4$ with respect to K_Y [Ha, Theorem 7.4]. Therefore P has to be cA_2 .

We take an isomorphism $P \in X \cong o \in (x_1x_2 + g_3(x_3, x_4) + g_{\geq 4}(x_3, x_4) = 0) \subset \mathbb{C}^4$, where g_3 is the part of degree 3 and $g_{\geq 4}$ is the part of degree greater than or equal to 4. To see (i) we have to show that g_3 is cubic, but this comes from the property that there exists a hyperplane section of $P \in X$ which is Du Val of type E_6 at P (6.9). $\square$

Example 7.10. There exists an example of (7.9). The weighted blowup of $o \in (x_1^2 + x_2^2 + x_3^3 + x_1x_4^2 = 0) \subset \mathbb{C}^4$ with weights $\text{wt}(x_1, x_2, x_3, x_4) = (4, 3, 2, 1)$ is such an example.

We have (7.4), (7.5), (7.8) and (7.9). We can easily check the sufficient condition for f to be a divisorial contraction, and obtain the classification (2.12).

References

- [Ar1] M. Artin, On the solutions of analytic equations, *Invent. Math.* **5**, (1968) 277-291
- [Ar2] M. Artin, Algebraic approximation of structures over complete local rings, *Publ. Math. Inst. Hautes Étud. Sci.* **36**, (1969) 23-58
- [Co] A. Corti, Singularities of linear systems and 3-fold birational geometry, *Explicit birational geometry of 3-folds*, *Lond. Math. Soc. Lect. Note Ser.* **281**, (2000) 259-312
- [CM] A. Corti and M. Mella, Birational geometry of terminal quartic 3-folds I, preprint (2000)
- [Ha] T. Hayakawa, Blowing ups of 3-dimensional terminal singularities, *Publ. Res. Ins. Math. Sci.* **35**, (1999) 515-570
- [Ka1] M. Kawakita, Divisorial contractions in dimension three which contract divisors to smooth points, *Invent. Math.* **145**, (2001) 105-119
- [Ka2] M. Kawakita, Divisorial contractions in dimension three which contract divisors to compound A_1 points, to appear in *Compos. Math.*
- [Km] Y. Kawamata, Divisorial contractions to 3-dimensional terminal quotient singularities, *Higher-dimensional complex varieties*, de Gruyter, (1996) 241-246

- [KMM] Y. Kawamata, K. Matsuda and K. Matsuki, Introduction to the minimal model problem, *Adv. Stud. Pure Math.* **10**, (1987) 283-360
- [KM1] J. Kollár and S. Mori, Classification of three-dimensional flips, *J. Am. Math. Soc.* **5**, (1992) 533-703
- [KM2] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, *Camb. Tracts Math.* **134**, (1998)
- [KS] J. Kollár and N. Shepherd-Barron, Threefolds and deformations of surface singularities, *Invent. Math.* **91**, (1988) 299-338
- [Ma] D. Markushevich, Minimal discrepancy for a terminal cDV singularity is 1, *J. Math. Sci. Tokyo* **3**, (1996) 445-456
- [Mo1] S. Mori, On 3-dimensional terminal singularities, *Nagoya Math. J.* **98**, (1985) 43-66
- [Mo2] S. Mori, Flip theorem and the existence of minimal models for 3-folds, *J. Am. Math. Soc.* **1**, (1988) 117-253
- [Re1] M. Reid, Minimal models of canonical 3-folds, *Adv. Stud. Pure Math.* **1**, (1983) 131-180
- [Re2] M. Reid, Projective morphisms according to Kawamata, preprint (1983)
- [Re3] M. Reid, Young person's guide to canonical singularities, *Proc. Symp. Pure Math.* **46**, (1987) 345-414
- [Sh] V. Shokurov, Smoothness of a general anticanonical divisor on a Fano variety, *Izv. Akad. Nauk SSSR Ser. Mat.* **43**, (1979) 430-441
- [Ta] H. Takagi, On classification of $\mathbb{Q}$ -Fano 3-folds of Gorenstein index 2 II, preprint (2000)

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO,
 3-8-1 KOMABA, MEGURO, TOKYO 153-8914, JAPAN
email address: kawakita@ms.u-tokyo.ac.jp